

North Carolina, 2010

Primary and secondary syphilis

- CASES: North Carolina reported 396 cases in 2010.
- RATES: North Carolina ranked 18 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 4.2 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In North Carolina, the rate among males was 7.4 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 1.1 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In North Carolina, the race/ethnicity adjusted rates per 100,000 population were 0.9 among whites, 15.8 among blacks, 1.7 among Hispanics, 0.5 among Asian/Pacific Islanders, and 0.9 among American Indians/Alaska Natives.
- The rate among blacks was 17.6 times that of whites.
- The rate among Hispanics was 1.9 times that of whites.


Congenital syphilis

- CASES: North Carolina reported 10 cases in 2010.
- RATES: North Carolina ranked 15 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 7.6 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, North Carolina, 2001-2010



Total P&S Syphilis Rates by County, North Carolina, 2010



Rate (per 100,000 population)

