

New York, 2010

Primary and secondary syphilis

- CASES: New York reported 1,098 cases in 2010.
- RATES: New York ranked 10 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 5.6 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In New York, the rate among males was 11.1 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.5 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In New York, the race/ethnicity adjusted rates per 100,000 population were 2.5 among whites, 14.7 among blacks, 8.9 among Hispanics, 1.1 among Asian/Pacific Islanders, and 2.7 among American Indians/Alaska Natives.
- The rate among blacks was 5.9 times that of whites.
- The rate among Hispanics was 3.6 times that of whites.
- The rate among American Indians/Alaska Natives was 1.1 times that of whites.

Congenital syphilis

- CASES: New York reported 16 cases in 2010.
- RATES: New York ranked 18 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 6.3 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, New York, 2001-2010



Total P&S Syphilis Rates by County, New York, 2010

