

Ohio, 2010

Primary and secondary syphilis

- CASES: Ohio reported 528 cases in 2010.
- RATES: Ohio ranked 15 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 4.6 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Ohio, the rate among males was 7.0 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 2.2 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Ohio, the race/ethnicity adjusted rates per 100,000 population were 1.7 among whites, 24.5 among blacks, 2.5 among Hispanics, 0.5 among Asian/Pacific Islanders, and 3.3 among American Indians/Alaska Natives.
- The rate among blacks was 14.4 times that of whites.
- The rate among Hispanics was 1.5 times that of whites.
- The rate among American Indians/Alaska Natives was 1.9 times that of whites.

Congenital syphilis

- CASES: Ohio reported 10 cases in 2010.
- RATES: Ohio ranked 17 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 6.6 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Ohio, 2001-2010



Total P&S Syphilis Rates by County, Ohio, 2010

