

# South Carolina, 2010

## Primary and secondary syphilis

- CASES: South Carolina reported 155 cases in 2010.
- RATES: South Carolina ranked 23 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 3.4 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In South Carolina, the rate among males was 6.6 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.4 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In South Carolina, the race/ethnicity adjusted rates per 100,000 population were 0.8 among whites, 9.7 among blacks, 1.5 among Hispanics, 1.5 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 12.1 times that of whites.
- The rate among Hispanics was 1.9 times that of whites.
- The rate among Asian/Pacific Islanders was 1.9 times that of whites.


## Congenital syphilis

- CASES: South Carolina reported 0 cases in 2010.
- RATES: South Carolina had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, South Carolina, 2001-2010



## Total P&S Syphilis Rates by County, South Carolina, 2010



Rate (per 100,000 population)

