

Oklahoma, 2011

Primary and secondary syphilis

- CASES: Oklahoma reported 84 cases in 2011.
- RATES: Oklahoma ranked 36 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.2 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Oklahoma, the rate among males was 3.9 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.6 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Oklahoma, the race/ethnicity adjusted rates per 100,000 population were 1.6 among whites, 8.9 among blacks, 1.2 among Hispanics, 4.0 among Asian/Pacific Islanders, and 1.4 among American Indians/Alaska Natives.
- The rate among blacks was 5.6 times that of whites.
- The rate among Asian/Pacific Islanders was 2.5 times that of whites.


Congenital syphilis

- CASES: Oklahoma reported 2 cases in 2011.
- RATES: Oklahoma ranked 21 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 3.7 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Oklahoma, 2002 – 2011



Total P&S Syphilis Rates by County, Oklahoma, 2011



Rate (per 100,000 population)

