# Ohio, 2012

## Primary and secondary syphilis

- CASES: Ohio reported 425 cases in 2012.
- RATES: Ohio ranked 23 among 50 states reporting at least one case of P&S syphilis with 3.7 cases per 100,000 population compared to the U.S. rate of 5.0.

## Rates of primary and secondary syphilis, by sex

- In Ohio, the rate among males was 6.0 cases per 100,000 population compared to the U.S. male rate of 9.3.
- The rate among females was 1.4 compared to the U.S. female rate of 0.9.

## Rates of primary and secondary syphilis, by race/ethnicity

- In Ohio, the race/ethnicity adjusted rates per 100,000 population were 1.5 among whites, 17.3 among blacks, 3.5 among Hispanics, 0.5 among Asians, 0.0 among Native Hawaiians and Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 11.4 times that of whites.
- The rate among Hispanics was 2.3 times that of whites.

## Congenital syphilis

- CASES: Ohio reported 16 cases in 2012.
- RATES: Ohio ranked 8 among among 30 states reporting at least one case of congenital syphilis with a rate of 11.0 cases per 100,000 live births compared to the U.S. rate of 7.8.

**Syphilis: P&S Rates and Congenital Cases, Ohio, 2003–2012**




**Total P&S Syphilis Rates by County, Ohio, 2012**