Tripler Army Medical Center, Hawaii – 2002 (N=26)

Figure A. Age of GISP participants, in years, 2002



Figure B. Race/ethnicity of GISP participants, 2002



Figure C. Percentage of GISP participants identifying as men who have sex with men, 1988–2002

Data not collected

Figure D. Reason for visit among GISP participants, 2002



Tripler Army Medical Center, Hawaii – 2002 (N=26)

Figure E. Previous episode of gonorrhea among GISP participants, 1991–2002

Data not available

Figure F. Drugs used to treat gonorrhea among GISP participants, 2002



Figure G. Drugs used to treat *Chlamydia trachomatis* infection among GISP participants, 2002



Figure H. Resistance to penicillin and tetracycline among GISP isolates, 2002



Tripler Army Medical Center, Hawaii – 2002 (N=26)

Figure I. Decreased susceptibility to ceftriaxone among GISP isolates, 1988–2002

No isolates with decreased susceptibility to ceftriaxone have been identified at this clinic.

Figure J. Decreased susceptibility to cefixime among GISP isolates, 1992–2002

No isolates with decreased susceptibility to cefixime have been identified at this clinic.

Figure K. Intermediate resistance and resistance to ciprofloxacin among GISP isolates, 1990–2002

No isolates with intermediate resistance or resistance to ciprofloxacin have been identified at this clinic.

Figure L. Decreased susceptibility to azithromycin among GISP isolates, 1992–2002

No isolates with decreased susceptibility to azithromycin have been identified at this clinic.