



Tripler Army Medical Center, Hawaii (N=6)

Figure A. Age of GISP participants, in years, 2009



Tripler Army Medical Center, Hawaii (N=6)

Figure B. Race/ethnicity of GISP participants, 2009




## Tripler Army Medical Center, Hawaii

Figure C. Percentage of GISP participants identifying as men who have sex with men, 2000-2009

Data not collected


Tripler Army Medical Center, Hawaii (N=6)

Figure D. Drugs used to treat gonorrhea among GISP participants, 2009




Tripler Army Medical Center, Hawaii (N=6)

Figure E. Drugs used to treat *Chlamydia trachomatis* infection among GISP participants, 2009




Tripler Army Medical Center, Hawaii (N=6)

Figure F. Resistance to penicillin, tetracycline, and ciprofloxacin among GISP isolates, 2009




# Tripler Army Medical Center, Hawaii

Figure G. Distribution of Minimum Inhibitory Concentrations (MICs) to ceftriaxone among GISP isolates, 2005-2009



# Tripler Army Medical Center, Hawaii


Figure H. Distribution of Minimum Inhibitory Concentrations (MICs) to cefixime among GISP isolates, 2005-2006 and 2009



NOTE: Isolates were not tested for cefixime susceptibility in 2007 and 2008.

# Tripler Army Medical Center, Hawaii

Figure I. Intermediate resistance and resistance to ciprofloxacin among GISP isolates, 1990-2009



Note: Susceptibility to ciprofloxacin first measured in 1990.

# Tripler Army Medical Center, Hawaii

Figure J. Distribution of Minimum Inhibitory Concentrations (MICs) to azithromycin among GISP isolates, 2005-2009

