



Columbus, Ohio (N=113)

Figure A. Age of GISP participants, in years, 2012




Columbus, Ohio (N=113)

Figure B. Race/ethnicity of GISP participants, 2012



# Columbus, Ohio


Figure C. Percentage of GISP participants identifying as men who have sex with men, 2000-2012



\*Note: Site participation in GISP began in 2012.


Columbus, Ohio (N=113)

Figure D. Primary antibiotic used to treat gonorrhea among GISP participants, 2012




Columbus, Ohio (N=113)

Figure E. Secondary antibiotic used to treat gonorrhea among GISP participants, 2012




Columbus, Ohio (N=113)

Figure F. Resistance to penicillin, tetracycline, and ciprofloxacin among GISP isolates, 2012




# Columbus, Ohio

Figure G. Distribution of Minimum Inhibitory Concentrations (MICs) of ceftriaxone among GISP isolates, 2008-2012



# Columbus, Ohio


Figure H. Distribution of Minimum Inhibitory Concentrations (MICs) of cefixime among GISP isolates, 2009-2012



NOTE: Isolates were not tested for cefixime susceptibility in 2007 and 2008.

# Columbus, Ohio

Figure I. Intermediate resistance and resistance to ciprofloxacin among GISP isolates, 2000-2012



Note: Susceptibility to ciprofloxacin first measured in 1990. Site participation in GISP began in 2012.

# Columbus, Ohio

Figure J. Distribution of Minimum Inhibitory Concentrations (MICs) of azithromycin among GISP isolates, 2008-2012

