




Syphilis Surveillance Supplemental Slides

2015–2019

July 2021

Reported* Injection Drug Use, Methamphetamine Use, Heroin Use, and Sex with a PWID Among Primary and Secondary Syphilis Cases, United States, 2015–2019




* Proportion reporting injection drug use, methamphetamine use, heroin use, or sex with a PWID within the last 12 months calculated among cases with known data (cases with missing or unknown responses were excluded from the denominator).

Abbreviations: PWID = person who injects drugs



Reported* Injection Drug Use, Methamphetamine Use, Heroin Use, and Sex with a PWID Among MSM with Primary and Secondary Syphilis, United States, 2015–2019




* Proportion reporting injection drug use, methamphetamine use, heroin use, or sex with a PWID within the last 12 months calculated among cases with known data (cases with missing or unknown responses were excluded from the denominator).

Abbreviations: PWID = person who injects drugs; MSM = gay, bisexual, or other men who have sex with men



Reported* Injection Drug Use, Methamphetamine Use, Heroin Use, and Sex with a PWID Among MSW with Primary and Secondary Syphilis, United States, 2015–2019




* Proportion reporting injection drug use, methamphetamine use, heroin use, or sex with a PWID within the last 12 months calculated among cases with known data (cases with missing or unknown responses were excluded from the denominator).

Abbreviations: PWID = person who injects drugs; MSW = men who have sex with women only



Reported* Injection Drug Use, Methamphetamine Use, Heroin Use, and Sex with a PWID Among Women with Primary and Secondary Syphilis, United States, 2015–2019




* Proportion reporting injection drug use, methamphetamine use, heroin use, or sex with a PWID within the last 12 months calculated among cases with known data (cases with missing or unknown responses were excluded from the denominator).

Abbreviations: PWID = person who injects drugs



Injection Drug Use

Reported Injection Drug Use* Among Primary and Secondary Syphilis Cases by Sex and Sex of Sex Partners, United States, 2015–2019




* Proportion reporting injection drug use within the last 12 months calculated among cases with known data (cases with missing or unknown responses were excluded from the denominator).

Abbreviations: MSM = gay, bisexual, or other men who have sex with men; MSW = men who have sex with women only



Reported Injection Drug Use* Among MSM with Primary and Secondary Syphilis by Region, United States, 2015–2019




* Proportion reporting injection drug use within the last 12 months calculated among cases with known data (cases with missing or unknown responses were excluded from the denominator).

Abbreviations: MSM = gay, bisexual, or other men who have sex with men



Reported Injection Drug Use* Among MSW with Primary and Secondary Syphilis by Region, United States, 2015–2019




* Proportion reporting injection drug use within the last 12 months calculated among cases with known data (cases with missing or unknown responses were excluded from the denominator).

Abbreviations: MSW = men who have sex with women only



Reported Injection Drug Use* Among Women with Primary and Secondary Syphilis by Region, United States, 2015–2019




* Proportion reporting injection drug use within the last 12 months calculated among cases with known data (cases with missing or unknown responses were excluded from the denominator).



Methamphetamine Use

Reported Methamphetamine Use* Among Primary and Secondary Syphilis Cases by Sex and Sex of Sex Partners, United States, 2015–2019




* Proportion reporting methamphetamine use within the last 12 months calculated among cases with known data (cases with missing or unknown responses were excluded from the denominator).

Abbreviations: MSM = gay, bisexual, or other men who have sex with men; MSW = men who have sex with women only



Reported Methamphetamine Use* Among MSM with Primary and Secondary Syphilis by Region, United States, 2015–2019




* Proportion reporting methamphetamine use within the last 12 months calculated among cases with known data (cases with missing or unknown responses were excluded from the denominator).

Abbreviations: MSM = gay, bisexual, or other men who have sex with men



Reported Methamphetamine Use* Among MSW with Primary and Secondary Syphilis by Region, United States, 2015–2019




* Proportion reporting methamphetamine use within the last 12 months calculated among cases with known data (cases with missing or unknown responses were excluded from the denominator).

Abbreviations: MSW = men who have sex with women only



Reported Methamphetamine Use* Among Women with Primary and Secondary Syphilis by Region, United States, 2015–2019




* Proportion reporting methamphetamine use within the last 12 months calculated among cases with known data (cases with missing or unknown responses were excluded from the denominator)



Heroin Use

Reported Heroin Use* Among Primary and Secondary Syphilis Cases by Sex and Sex of Sex Partners, United States, 2015–2019




* Proportion reporting heroin use within the last 12 months calculated among cases with known data (cases with missing or unknown responses were excluded from the denominator).

Abbreviations: MSM = gay, bisexual, or other men who have sex with men; MSW = men who have sex with women only



Reported Heroin Use* Among MSM with Primary and Secondary Syphilis by Region, United States, 2015–2019




* Proportion reporting heroin use within the last 12 months calculated among cases with known data (cases with missing or unknown responses were excluded from the denominator).

Abbreviations: MSM = gay, bisexual, or other men who have sex with men



Reported Heroin Use* Among MSW with Primary and Secondary Syphilis by Region, United States, 2015–2019




* Proportion reporting heroin use within the last 12 months calculated among cases with known data (cases with missing or unknown responses were excluded from the denominator).

Abbreviations: MSW = men who have sex with women only



Reported Heroin Use* Among Women with Primary and Secondary Syphilis by Region, United States, 2015–2019




* Proportion reporting heroin use within the last 12 months calculated among cases with known data (cases with missing or unknown responses were excluded from the denominator).



Sex with a Person Who Injects Drugs

Reported Sex with a PWID* Among Primary and Secondary Syphilis Cases by Sex and Sex of Sex Partners, United States, 2015–2019




* Proportion reporting sex with a PWID within the last 12 months calculated among cases with known data (cases with missing or unknown responses were excluded from the denominator).

Abbreviations: PWID = person who injects drugs; MSM = gay, bisexual, or other men who have sex with men; MSW = men who have sex with women only



Reported Sex with a PWID* Among MSM with Primary and Secondary Syphilis by Region, United States, 2015–2019




* Proportion reporting sex with a PWID within the last 12 months calculated among cases with known data (cases with missing or unknown responses were excluded from the denominator).

Abbreviations: PWID = person who injects drugs; MSM = gay, bisexual, or other men who have sex with men



Reported Sex with a PWID* Among MSW with Primary and Secondary Syphilis by Region, United States, 2015–2019




* Proportion reporting sex with a PWID within the last 12 months calculated among cases with known data (cases with missing or unknown responses were excluded from the denominator).

Abbreviations: PWID = person who injects drugs; MSW = men who have sex with women only



Reported Sex with a PWID* Among Women with Primary and Secondary Syphilis by Region, United States, 2015–2019



* Proportion reporting sex with a PWID within the last 12 months calculated among cases with known data (cases with missing or unknown responses were excluded from the denominator).

Abbreviations: PWID = person who injects drugs

