

APPENDIX

APPENDIX

Interpreting STD Surveillance Data

Sexually Transmitted Disease Surveillance, 2007 presents surveillance information derived from the official statistics for the reported occurrence of nationally notifiable sexually transmitted diseases in the United States, test positivity and prevalence data from numerous prevalence monitoring initiatives, sentinel surveillance of gonococcal antimicrobial resistance, and national health care services surveys.

Nationally Notifiable STD Surveillance

Nationally notifiable STD surveillance data are collected and compiled from reports sent by the STD control programs and health departments in the 50 states, the District of Columbia, selected cities, U.S. dependencies and possessions, and independent nations in free association with the United States to the Division of STD Prevention (DSTDP), National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention (NCHHSTP), Centers for Disease Control and Prevention (CDC). Included among the dependencies, possessions, and independent nations are Guam, Puerto Rico, and the Virgin Islands. These entities are identified as “outlying areas” of the United States in selected figures and tables.

Reporting Formats

STD morbidity data presented in this report are compiled from a combination of data reported on standardized hardcopy report forms and electronic data received via the National Electronic Telecommunications System for Surveillance (NETSS).

Summary Report Forms (hardcopy format)

The following hardcopy forms were used to report national STD morbidity data:

1. FORM CDC 73.998: *Monthly Surveillance Report of Early Syphilis*. This monthly hardcopy reporting form was used from 1984 to 2002 to report summary data for P&S syphilis and early latent syphilis by county and state.

2. FORM CDC 73.688: *Sexually Transmitted Disease Morbidity Report*. This quarterly hardcopy reporting form was used from 1963 to 2002 to report summary data for all stages of syphilis, congenital syphilis, gonorrhea, chancroid, chlamydia, and other STDs by sex and source of report (private vs. public) for the 50 states, Washington, D.C., and 64 selected cities (including San Juan, Puerto Rico) and outlying areas of the United States. Note: genital chlamydial infection became a nationally notifiable condition in 1996, and the form was modified to support reporting of chlamydia that year. Congenital syphilis was dropped from this aggregate form in 1995 and replaced by the case-specific CDC 73.126 form described below.

3. FORM CDC 73.2638: *Report of Civilian Cases of Primary & Secondary Syphilis, Gonorrhea, and Chlamydia by Reporting Source, Sex, Race/Ethnicity, and Age Group*. This annual hardcopy form was used from 1981 to 2002 to report summary data for P&S syphilis, gonorrhea, and chlamydia by age, race, sex and source (public vs. private) for all states and seven large cities (Baltimore, Chicago, New York City, Los Angeles, Philadelphia, San

Francisco, and Washington, D.C.), and outlying areas of the United States. Note: genital chlamydial infection became a nationally notifiable condition in 1996 and the form was modified to support reporting of chlamydia that year.

4. FORM CDC 73.126: *Congenital Syphilis (CS) Case Investigation and Report*. This case-specific hardcopy form was first used in 1983 and continues to be used to report detailed case-specific data for congenital syphilis in some reporting areas.

National Electronic Telecommunications System for Surveillance (NETSS, electronic format)

Notifiable STD data reported electronically through NETSS comprise the nationally notifiable disease information that is published in the *Morbidity and Mortality Weekly Report (MMWR)*.

As of December 31, 2003, all 50 states and Washington, D.C. had converted from summary hardcopy reporting to electronic submission of line-listed (i.e., case-specific) STD data via NETSS (41 reporting areas are submitting congenital syphilis surveillance data via NETSS). Puerto Rico converted to electronic reporting in 2006. Guam and the Virgin Islands continue to report using summary hardcopy forms.

Jurisdictions differ in their ability to resolve differences in total cases derived from summary hardcopy monthly, quarterly, and annual reports (as well as electronically submitted line-listed data). Thus, depending on the database used, there may be discrepancies in the total number of cases among the figures and tables for earlier years. In most instances, these discrepancies are less than 5% of total reported cases and have minimal impact on national case totals and rates. However, for a specific jurisdiction, the discrepancies may be larger.

Surveillance data and updates sent to CDC on hardcopy forms and for NETSS electronic data through June 25, 2008 have been included in this report. Data received after this date will appear in subsequent annual Surveillance Reports. The data presented in the figures and tables in this document supersede those in all earlier publications.

Population Denominators and Rate Calculations

2000—2007 Rates and Population

The National Center for Health Statistics released bridged race population counts for 2000–2006 resident population based on the Census 2000 counts. These estimates resulted from bridging the 31 race categories used in Census 2000, as specified in the 1997 Office of Management and Budget (OMB) standards, to the five race/ethnicity groups specified under the 1977 OMB standards.

Population estimates for Guam, Puerto Rico, and the Virgin Islands were obtained from the Bureau of Census web site: <http://www.census.gov/ipc/www/idb/tables.html>. The 2006–2007 rates for outlying areas were calculated using the 2006 population estimates.

Due to use of the updated population data, rates for the period 2000–2006 may be different from prior Surveillance Reports.

1990—1999 Rates and Population

The population counts for 1990–1999 incorporated the bridged single-race estimates of the April 1, 2000 resident population. These files were prepared by the U.S. Census Bureau with support from the National Cancer Institute.

1981—1989 Rates and Population

For the United States, rates were calculated using Bureau of the Census population estimates for 1981 through 1989 (Bureau of the Census; United States Population Estimates by Age, Sex and Race: 1980–1989 [Series P-25, No. 1045]; Washington: U.S. Government Printing Office, 1990; and United States Population Estimates by Age, Sex and Race: 1989 [Series P-25, No. 1057]; Washington: U.S. Government Printing Office, 1990.

1941—1980 Rates and Population

Rates for 1941–1980 are based on population estimates from the Bureau of Census and currently maintained by the Division of STD Prevention.

1941—2007 Congenital Syphilis Rates and Live Births

Congenital syphilis (CS) data in Table 1 of *Sexually Transmitted Disease Surveillance 2007* represent the number of congenital syphilis cases per 100,000 live births for all years during the period 1941 through 2007. Previous publications presented congenital syphilis rates per 100,000 population for 1941–1994 and rates for cases diagnosed at < 1 year of age per 100,000 live births for 1995–2005. To allow for trends in CS rates to be compared over the period 1941–2007, live births are used as the denominator for CS and case counts are no longer limited to those diagnosed within the first year of life. CS morbidity (i.e., case reports) is assigned by year of birth. Rates of congenital syphilis for 1963–1988 were calculated using published live birth data (NCHS; Vital Statistics Report, United States, 1988 [Vol. 1-Natality]). Congenital syphilis rates for 1989–2003 were calculated using live birth data based on information coded by the States and provided to the National Center for Health Statistics (NCHS) through the Vital Statistics Cooperative

Program. Rates for 2004–2007 were calculated using live birth data for 2004.

Reporting Practices

Although most areas generally adhere to the national notifiable STD case definitions collaboratively developed by the Council of State and Territorial Epidemiologists (CSTE) and CDC, there may be differences in the policies and systems for collecting surveillance data. Thus, comparisons of case numbers and rates between jurisdictions should be interpreted with caution. However, since case definitions and surveillance activities within a given area remain relatively stable over time, trends should be minimally affected by these differences. In many areas, the reporting from publicly supported institutions (e.g., STD clinics) has been more complete than from other sources (e.g., private practitioners). Thus, trends may not be representative of all segments of the population.

Reporting of Metropolitan Statistical Area-specific Surveillance Data

Sexually Transmitted Disease Surveillance, 2007 continues the presentation of STD incidence data and rates for the fifty Metropolitan Statistical Areas (MSAs) with the largest populations based on 2000 U.S. Census data. *Sexually Transmitted Disease Surveillance* reports prior to 2005 presented data by selected cities which estimated city-specific morbidity and were derived from county data. Since county data were used to estimate city-specific morbidity and current STD project areas' reporting practices do not support direct identification of city-specific morbidity reports, MSAs (described below) were chosen as a geographic unit smaller than a state or territory for presentation of STD morbidity data.

Metropolitan Statistical Areas are defined by the Office of Management and Budget

to provide nationally consistent definitions for collecting, tabulating, and publishing federal statistics for a set of geographic areas.¹ An MSA is associated with at least one urbanized area that has a population of at least 50,000. The MSA comprises the central county or counties containing the core, plus adjacent outlying counties having a high degree of social and economic integration with the central county as measured through commuting. The title of an MSA includes the name of its principal city with the largest Census 2000 population. If there are multiple principal cities, the names of the second largest and third largest principal cities appear in the title in order of descending population size.

The MSA concept has been used as a statistical representation of the social and economic linkages between urban cores and outlying, integrated areas. However, MSAs do not equate to an urban-rural classification; all counties included in MSAs and many other counties contain both urban and rural territory and populations. Programs that treat all parts of an MSA as if they were as urban as the densely settled core ignore the rural conditions that may exist in some parts of the area. In short, MSAs are not designed as a general purpose geographic framework for nonstatistical activities or for use in program funding formulas.

For more information on MSA definitions used in this report, please visit this web site: <http://www.census.gov/population/estimates/metro-city/03mfips.txt>

Management of Unknown, Missing or Invalid Age Group, Race/Ethnicity, and Sex Data

The percentage of unknown, missing or invalid data for age group, race/ethnicity, and sex varies from year to year, state to state, and by disease for reported STDs (Table A1).

When the percentage of unknown, missing, or invalid data for the variables — age group, race/ethnicity, and sex — exceeds 50% for any state, the state's incidence data and population data are excluded from the tables presenting data stratified by one or more of these variables. For those states reporting > 50% valid data for these variables, unknown, missing or invalid data are redistributed based on the state's distribution of known age group, race/ethnicity, and sex data, respectively. As a result of this procedure, incidence and rate data stratified by one or more of the variables—age group, race/ethnicity, and sex—may not accurately reflect total national incidence or rates.

Classification of STD Morbidity Reporting Sources

Prior to 1996, states classified the source of case reports as either private source (including private physicians, and private hospitals and institutions) or public (clinic) source (primarily STD clinics). As states began reporting morbidity data electronically in 1996, the classification categories for source of case reports expanded to include the following data sources: STD clinics, HIV counseling and testing sites, drug treatment clinics, family planning clinics, prenatal/obstetrics clinics, tuberculosis clinics, private physicians/HMOs, hospitals (inpatient), emergency rooms, correctional facilities, laboratories, blood banks, National Job Training Program, school-based clinics, mental health providers, military, Indian Health Service, and other unspecified sources. Analysis of the data reported electronically after 1996 confirmed that the new STD clinic source of report data corresponded to the earlier reporting source category, public (clinic) source. Therefore, source of case report data for the period 1984–2007 are presented as STD clinic or non-STD clinic only (Table A2).

Definition of DHHS Regions

The ten U.S. Department of Health and Human Services (DHHS) regions referred to in the text and figures include the following jurisdictions: Region I = Connecticut, Maine, Massachusetts, New Hampshire, Rhode Island, and Vermont; Region II = New Jersey, New York, Puerto Rico, and U.S. Virgin Islands; Region III = Delaware, District of Columbia, Maryland, Pennsylvania, Virginia, and West Virginia; Region IV = Alabama, Florida, Georgia, Kentucky, Mississippi, North Carolina, South Carolina, and Tennessee; Region V = Illinois, Indiana, Michigan, Minnesota, Ohio, and Wisconsin; Region VI = Arkansas, Louisiana, New Mexico, Oklahoma, and Texas; Region VII = Iowa, Kansas, Missouri, and Nebraska; Region VIII = Colorado, Montana, North Dakota, South Dakota, Utah, and Wyoming; Region IX = Arizona, California, Guam, Hawaii, and Nevada; and Region X = Alaska, Idaho, Oregon, and Washington.

Chlamydia Morbidity Reporting

Trends in chlamydia morbidity reporting from many areas are more reflective of changes in diagnosis, screening, and reporting practices than of actual trends in disease incidence. In particular, morbidity trends are likely highly influenced by changes in test technology, as areas increase their usage of more sensitive nucleic acid amplification tests. As areas develop chlamydia prevention and control programs, including improved surveillance systems to monitor trends, the data should improve and become more representative of true trends in disease.

Syphilis Morbidity Reporting

“Total syphilis” or “all stages of syphilis” includes primary, secondary, latent (including early latent, late latent, and latent syphilis of unknown duration), neurosyphilis, late (including late syphilis

with clinical manifestations other than neurosyphilis), and congenital syphilis.

In 1996, the syphilis stage, “late syphilis with clinical manifestations other than neurosyphilis (late benign and cardiovascular syphilis)”, was added to the syphilis case definition (see STD Case Definitions in this **Appendix**).

While neurosyphilis can occur at almost any stage of syphilis, between 1996 and 2005, it was classified and reported as one of several mutually exclusive stages of syphilis. Beginning in 2005, neurosyphilis was no longer classified or reported as a distinct stage of syphilis. The Division of STD Prevention is reviewing these data collection and reporting changes to determine their impact on syphilis surveillance case definitions.

Congenital Syphilis Morbidity Reporting

In 1988, the surveillance case definition for congenital syphilis was changed. This case definition has greater sensitivity than the former definition.² In addition, many areas have greatly enhanced active case finding for congenital syphilis since 1988. For these reasons, the number of reported cases increased dramatically during 1989–1991. All reporting areas had implemented the new case definition for reporting congenital syphilis by January 1, 1992.

In addition to changing the case definition for congenital syphilis, CDC introduced a new data collection form (CDC 73.126) in 1990 (revised October 2003). Since 1995, the data collected on this form have been used for reporting congenital syphilis cases and associated rates. This form is used to collect individual case information which allows more thorough analysis of case characteristics. For the purpose of analyzing race/ethnicity, cases are classified by race/ethnicity of the mother. Congenital syphilis cases were reported by state and

city of residence of the mother for the period 1995 through 2007.

Congenital syphilis reporting may be delayed as a result of case investigation and validation. Congenital syphilis cases for prior years are added to CDC's surveillance databases throughout the year. Congenital syphilis data reported after publication of the current annual Surveillance Report will appear in subsequent surveillance reports and are assigned by case patient's year of birth.

Chlamydia, Gonorrhea, and Syphilis Prevalence Monitoring

Chlamydia and gonorrhea test positivity and syphilis seroreactivity were calculated for the following: women attending family planning clinics and prenatal clinics, men and women entering the National Job Training Program, women attending Indian Health Service clinics, men attending STD clinics and primary care clinics participating in the MSM Prevalence Monitoring Project, and men and women entering corrections facilities. Positivity was calculated by dividing the number of positive tests for chlamydia, gonorrhea, or syphilis (numerator) by the total number of positive and negative tests for each disease (denominator) and was expressed as a percentage. Except for the National Job Training Program screening data, these data sources may include more than one test from the same individual if that person was tested more than once during a year.

To increase the stability of the annual National Job Training Program prevalence estimates, chlamydia or gonorrhea prevalence data are presented when valid test results for 100 or more students per year are available for the population subgroup and state. The majority of the National Job Training Program's chlamydia screening tests are tested by a single national contract laboratory, which provides those data to CDC. Gonorrhea

screening tests for male and female students in many training centers are tested by local laboratories; these data are not available to CDC. To insure that state-specific gonorrhea screening data presented here are representative of all students entering training centers, gonorrhea test results for students at centers submitting specimens to the national contract laboratory are included only if the number of gonorrhea tests submitted is greater than 90% of the number of chlamydia tests submitted from the same center for the same time period.

Various laboratory test methods were used for all of these data sources. No adjustments for laboratory test type and sensitivity were made to any figures presenting test positivity or prevalence data. For more details on chlamydia prevalence, refer to the following annual publication: Centers for Disease Control and Prevention. *Sexually Transmitted Disease Surveillance 2007 Supplement: Chlamydia Prevalence Monitoring Project Annual Report 2007*. Atlanta, GA: U.S. Department of Health and Human Services (available first quarter 2009).

In the MSM Prevalence Monitoring Project, the syphilis seroreactivity data in most instances do not reflect confirmatory testing and thus biologic false positive test results were not systematically excluded. The extent to which these data reflect prevalence of active syphilis infection varies by site. Similarly, in the Corrections Prevalence Monitoring Project, syphilis seroreactivity test results were not confirmed. Only a few juvenile corrections sites submitted data to CDC, making overall interpretation difficult due to the small sample size. Because only selected corrections facilities participated in the Corrections Prevalence Monitoring Project, state-specific positivity for syphilis, chlamydia, and gonorrhea may not be representative of all corrections facilities in the state.

Prevalence data for region- and state-specific figures were published with permission from the Regional Infertility Prevention Program, selected state STD prevention programs, the National Job Training Program, and the Indian Health Service.

Gonococcal Isolate Surveillance Project (GISP)

Data on antimicrobial susceptibility in *Neisseria gonorrhoeae* were collected through the Gonococcal Isolate Surveillance Project (GISP), a sentinel system of 30 STD clinics and five regional laboratories located throughout the United States. For more details on findings from GISP gonorrhea surveillance activities, refer to the following annual publication: Centers for Disease Control and Prevention. *Sexually Transmitted Disease Surveillance 2007 Supplement: Gonococcal Isolate Surveillance Project (GISP) Annual Report 2007* (available first quarter, 2009). Atlanta, GA: U.S. Department of Health and Human Services, 2008.

Other Surveillance Data Sources

National Health and Nutrition Examination Survey (NHANES)

NHANES is composed of a series of cross-sectional surveys designed to provide national statistics on the health and nutritional status of the general household population through household interviews, standardized physical examinations, and the collection of biological samples in special mobile examination centers. In 1999, NHANES became a continuous survey with data released every 2 years. The sampling plan of the survey is a stratified, multistage, probability cluster design that selects a sample representative of the U.S. civilian noninstitutionalized population.

National Disease and Therapeutic Index (NDTI)

The information on the number of initial visits to private physicians' offices for sexually transmitted diseases was based on analysis of data from the National Disease and Therapeutic Index (NDTI) (machine-readable files or summary statistics for the period 1966 through 2007). The NDTI is a probability sample survey of private physicians' clinical management practices. For more information on this database, contact IMS Health, 660 W. Germantown Pike, Plymouth Meeting, PA 19462; Telephone: (800) 523-5333.

National Hospital Discharge Survey (NHDS)

The information on patients hospitalized for pelvic inflammatory disease or ectopic pregnancy was based on analysis of data from the National Hospital Discharge Survey (machine-readable files for years 1980–2006), an ongoing nationwide sample survey of medical records of patients discharged from acute care hospitals in the United States, conducted by the National Center for Health Statistics. For more information, see *Graves EJ. 1988 Summary: National Hospital Discharge Survey; Advance data No. 185*; Hyattsville (MD): National Center for Health Statistics, 1990.

National Hospital Ambulatory Medical Care Survey (NHAMCS-ER)

The National Hospital Ambulatory Medical Care Survey (NHAMCS-ER) (machine-readable files for 1995–2006) was used to obtain estimates of the number of emergency room visits for pelvic inflammatory disease among women ages 15 to 44 years. The estimates generated using these data sources (NHDS and NHAMCS-ER) are based on statistical

surveys and therefore have sampling variability associated with the estimates.

Healthy People 2010 Objectives

Healthy People 2010³ (HP2010) is a set of health objectives for the U.S. to achieve over the first decade of the new century. It is used by people, States, communities, professional organizations, and others to help develop programs to improve health. HP2010 builds on initiatives pursued over the past two decades. The 1979 Surgeon General's Report, Healthy People, and Healthy People 2000: National Health Promotion and Disease Prevention Objectives established national health objectives and served as the basis for the development of State and community plans. Like its predecessors, Healthy People 2010 was developed through a broad consultation process, built on the best scientific knowledge and designed to measure programs over time. Healthy People 2010 is organized into 28 focus areas, each with objectives and measures designed to drive action that will support two overarching goals: 1) increasing the quality and years of healthy life and 2) eliminating health disparities.

Focus area 25 of Healthy People 2010 — Sexually Transmitted Diseases, — contains objectives and measures related to STDs. The baselines, HP2010 targets and annual progress toward the targets are reported in Table A3. The year 2010 targets for the diseases addressed in this report are: primary and secondary syphilis — 0.2 case

per 100,000 population; congenital syphilis — 1.0 case per 100,000 live births; and gonorrhea — 19.0 cases per 100,000 population. An additional target established in the HP2010 objectives is to reduce the *Chlamydia trachomatis* test positivity to 3% among females aged 15 to 24 years who attend family planning and STD clinics and among males aged 15 to 24 who attend STD clinics.

Government Performance and Results Act of 1993 (GPRA) Goals

The Government Performance and Results Act of 1993 (GPRA) was enacted by Congress to increase the confidence of citizens in the capability of the federal government, to increase the effectiveness and accountability of federal programs, to improve service delivery, to provide agencies a uniform tool for internal management and to assist Congressional decision making. GPRA requires each agency to have a performance plan with long-term outcomes and annual, measurable performance goals and to report on these plans annually, comparing results with annual goals. There are two STD GPRA goals: 1) reduction in pelvic inflammatory disease (PID) and 2) elimination of syphilis. Each of these goals has measures. The long-term goals and measures of progress are reported in Table A4.

¹ Office of Management and Budget. Standards for Defining Metropolitan and Micropolitan Statistical Areas: Notice Federal Register December 27, 2000; 65(249):82228-38.

² Kaufman RE, Jones OG, Blount JH, Wiesner PJ. Questionnaire survey of reported early congenital syphilis: problems in diagnosis, prevention, and treatment. *Sexually Transmitted Diseases* 1977;4:135-9.

³ U.S. Department of Health and Human Services. *Healthy People 2010*. 2nd ed. With Understanding and Improving Health and Objectives for Improving Health. 2 vols. Washington, DC: U.S. Government Printing Office, November 2000.

Table A1. Selected STDs — Percentage of unknown, missing, or invalid values for selected variables by state and by nationally notifiable STD, 2007

State	Primary and Secondary Syphilis				Gonorrhea			Chlamydia		
	Percent Unknown Race/Ethnicity	Percent Unknown Age	Percent Unknown Sex	Percent Unknown Sex of Partners	Percent Unknown Race/Ethnicity	Percent Unknown Age	Percent Unknown Sex	Percent Unknown Race/Ethnicity	Percent Unknown Age	Percent Unknown Sex
Alabama	1.8	0.0	0.0	4.5	24.7	0.3	0.0	28.0	0.3	0.0
Alaska*	0.0	0.0	0.0	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Arizona	22.6	0.0	0.7	80.1	27.9	0.0	0.0	34.8	0.0	0.1
Arkansas	2.5	0.0	0.0	40.2	5.2	0.4	0.1	5.9	0.2	0.0
California	2.4	0.1	0.1	9.7	33.0	0.5	0.4	33.7	0.5	0.4
Colorado	7.0	0.0	0.0	8.8	31.9	0.0	0.0	55.5	0.1	0.0
Connecticut	2.6	0.0	0.0	2.6	26.6	0.5	0.0	35.4	0.8	0.0
Delaware	5.6	0.0	0.0	100.0	1.8	0.0	0.0	3.1	0.0	0.0
District of Columbia	1.1	0.0	0.0	21.3	29.0	0.7	0.5	41.4	0.6	0.4
Florida	7.8	0.0	0.0	13.4	12.5	0.1	0.0	15.8	0.1	0.0
Georgia	2.4	0.1	0.1	20.0	37.1	0.9	0.6	43.9	0.9	0.6
Hawaii*	0.0	0.0	0.0	0.0	37.0	0.2	0.0	52.1	0.1	0.0
Idaho*	100.0	0.0	0.0	0.0	27.1	0.0	1.9	26.0	0.1	0.9
Illinois	4.1	0.0	0.0	8.4	8.9	0.0	0.0	12.3	0.0	0.0
Indiana	3.7	0.0	0.0	0.0	14.7	0.0	0.3	18.0	0.0	0.5
Iowa	4.8	0.0	0.0	9.5	7.5	0.1	0.0	11.8	0.2	0.0
Kansas	0.0	0.0	0.0	7.1	23.8	0.2	0.0	35.9	0.4	0.0
Kentucky	3.6	0.0	0.0	7.1	18.8	0.1	0.1	22.9	0.2	0.1
Louisiana	2.1	0.0	0.0	100.0	15.0	0.3	1.0	22.6	0.5	1.4
Maine*	0.0	0.0	0.0	11.1	7.6	0.8	0.0	21.1	0.4	0.1
Maryland	2.0	0.0	0.0	13.0	22.7	0.4	0.0	28.6	0.4	0.1
Massachusetts	7.1	0.0	0.0	9.7	31.0	0.7	0.0	40.1	0.8	0.1
Michigan	0.8	0.0	0.0	17.1	40.2	0.3	0.3	44.6	0.3	0.4
Minnesota	6.8	0.0	0.0	0.0	14.2	0.0	0.0	15.4	0.0	0.0
Mississippi	0.8	0.8	0.0	9.0	17.4	0.1	0.0	19.3	0.1	0.0
Missouri	0.0	0.0	0.0	2.5	15.9	0.1	0.0	21.7	0.1	0.0
Montana*	25.0	0.0	0.0	62.5	32.0	0.0	0.0	30.3	0.4	0.3
Nebraska*	25.0	0.0	0.0	0.0	23.3	0.0	0.3	28.7	0.1	0.3
Nevada	11.7	0.0	0.0	2.7	29.4	0.3	0.0	37.8	0.4	0.1
New Hampshire	0.0	0.0	0.0	3.3	8.0	0.0	0.0	14.3	0.0	0.0
New Jersey	0.9	0.0	0.0	17.6	34.0	0.6	0.0	47.4	0.7	0.1
New Mexico	10.9	0.0	0.0	4.3	16.0	0.2	0.0	11.8	0.3	0.0
New York	11.5	0.0	0.0	20.5	32.8	0.5	0.1	41.7	0.5	0.1
North Carolina	0.0	0.0	0.0	100.0	1.6	0.0	0.0	2.2	0.0	0.0
North Dakota*	0.0	0.0	0.0	0.0	10.3	0.0	0.0	11.1	0.3	0.1
Ohio	2.1	0.0	0.0	5.7	25.3	1.2	0.6	31.1	1.4	0.9
Oklahoma	0.0	0.0	0.0	15.4	1.6	0.0	0.0	2.2	0.0	0.0
Oregon	0.0	0.0	0.0	100.0	6.5	0.0	0.0	12.0	0.0	0.0
Pennsylvania	6.1	0.0	0.0	2.7	20.3	0.1	0.0	23.6	0.0	0.1
Rhode Island	2.8	0.0	0.0	0.0	11.2	0.0	0.2	20.3	0.2	0.1
South Carolina	0.0	0.0	0.0	3.3	26.7	0.6	0.2	29.3	0.3	0.3
South Dakota*	0.0	0.0	0.0	0.0	1.5	2.3	0.4	2.2	1.8	0.0
Tennessee	0.0	0.0	0.0	1.4	4.7	0.1	0.0	7.2	0.1	0.0
Texas	0.3	0.0	0.0	2.3	9.8	0.3	0.2	9.5	0.2	0.2
Utah	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0
Vermont	0.0	0.0	0.0	30.0	3.1	0.0	0.0	1.1	0.0	0.0
Virginia	0.0	0.0	0.0	3.5	6.0	0.1	0.1	11.8	0.1	0.0
Washington	8.4	0.0	0.0	23.4	21.1	1.1	0.0	20.7	1.2	0.0
West Virginia*	0.0	0.0	0.0	0.0	13.1	0.3	0.0	14.5	0.2	0.0
Wisconsin	0.0	0.0	0.0	4.5	26.1	0.0	0.2	28.7	0.0	0.1
Wyoming*	50.0	0.0	0.0	0.0	23.5	0.0	0.0	23.0	0.9	0.0
U.S. TOTAL	4.1	0.0	0.0	19.5	20.3	0.3	0.2	25.7	0.4	0.2

*Percentages for P&S syphilis are based on less than 10 cases.

NOTE: "Unknown" includes unknown, missing, or invalid data values.

Table A2. Reported cases of sexually transmitted disease by reporting source and sex: United States, 2007

<i>Disease</i>	<i>Non-STD Clinic</i>			<i>STD Clinic</i>			<i>Total*</i>		
	<i>Male</i>	<i>Female</i>	<i>Total</i>	<i>Male</i>	<i>Female</i>	<i>Total</i>	<i>Male</i>	<i>Female</i>	<i>Total</i>
Chlamydia	165,128	658,503	825,160	92,906	98,382	191,495	280,337	825,660	1,108,374
Gonorrhea	90,669	137,261	228,359	62,320	32,495	94,891	167,684	187,595	355,991
Primary Syphilis	1,742	177	1,919	1,018	120	1,138	2,837	303	3,140
Secondary Syphilis	4,860	841	5,703	1,834	497	2,334	6,932	1,389	8,326
Early Latent Syphilis	5,092	1,886	6,982	2,265	995	3,260	7,751	3,013	10,768
Late and Late Latent Syphilis†	8,646	4,927	13,582	2,375	1,378	3,755	11,637	6,604	18,256
Chancroid	5	8	13	1	2	3	8	15	23

*Totals include unknown sex and reporting source.

†Late and late latent syphilis includes late latent syphilis, latent syphilis of unknown duration, neurosyphilis, and late syphilis with clinical manifestations other than neurosyphilis.

See Appendix (Classification of STD Morbidity Reporting Sources).

Table A3. Healthy People 2010 Sexually Transmitted Diseases Objective Status

<i>HP 2010 Objectives</i>		<i>Baseline Year</i>	<i>Baseline</i>	<i>2003</i>	<i>2004</i>	<i>2005</i>	<i>2006</i>	<i>2007</i>	<i>HP 2010 Target</i>
25-1	Reduce the proportion of adolescents and young adults with <i>Chlamydia trachomatis</i> infections								
	a. Females aged 15 to 24 years attending family planning clinics	1997	5.0%	6.4%*	6.9%*	6.9%*	7.1%*	7.5%*	3.0%
	b. Females aged 15 to 24 years attending STD clinics	1997	12.2%	14.1%*	15.3%*	15.4%*	14.8%*	15.3%*	3.0%
	c. Males aged 15 to 24 years attending STD clinics	1997	15.7%	19.3%*	20.8%*	20.5%*	20.8%*	22.4%*	3.0%
	d. Females aged 24 years or less enrolled in National Job Training Program	2002	10.1%	9.9%	10.3%	9.2%	13.1%	13.2%	6.8%
25-2	Reduce the incidence of gonorrhea (new cases per 100,000 population)								
	a. Reduce gonorrhea incidence per 100,000 population	1997	123.0	115.2	112.4	114.6	119.7	118.9	19.0
	b. Females aged 15 to 44 years	2002	279.0	270.0	267.0	275.0	290.0	290.0	42.0
25-3	Eliminate sustained domestic transmission of primary and secondary syphilis (cases per 100,000 population)	1997	3.2	2.5	2.7	2.9	3.3	3.8	0.2
25-4	Reduce the proportion of adults aged 20 to 29 years with genital herpes infection	1988-94	17.0%	NA	10.0% [†]	NA	NA	NA	14.0%
25-6	Reduce the proportion of females aged 15 to 44 years who have ever required treatment for pelvic inflammatory disease (PID)	1995	8.0%	5.0% [†]	NA	NA	NA	NA	5.0%
25-7	Reduce the proportion of childless females with fertility problems who have had a sexually transmitted disease or who have required treatment for pelvic inflammatory disease (PID)	1995	27.0%	22.0% [†]	NA	NA	NA	NA	15.0%
25-9	Reduce congenital syphilis (cases per 100,000 live births)	1997	27.0	10.6	9.1	8.2	9.3	10.5	1.0
25-11	Adolescent sexual behavior (grades 9 through 12)								
	a. Increase the proportion of adolescents (grades 9 through 12) who abstain from sexual intercourse or use condoms if currently sexually active	1999	50.0	53.0	NA	53.0	NA	52.0	56.0
	b. Increase the proportion of adolescents (grades 9 through 12) who had sexual intercourse, but not in the past 3 months	1999	27.0	26.0	NA	27.0	NA	27.0	30.0
	c. Increase the proportion of adolescents (grades 9 through 12) who used condoms at last intercourse	1999	58.0	63.0	NA	63.0	NA	62.0	65.0
25-13	Increase the proportion of Tribal, state and local sexually transmitted disease programs that routinely offer hepatitis B vaccines to all STD clients	1998	5.0	NA	NA	NA	NA	NA	90.0
25-16	Increase the proportion of sexually active females aged 25 years and under who are screened annually for genital chlamydia infections								
	a. Females aged 25 years and under enrolled in commercial MCOs	2002	25.0	30.0	32.0	35.0	37.0	38.0	62.0
	b. Females aged 25 years and under enrolled in Medicaid MCOs	2002	41.0	45.0	47.0	51.0	52.0	52.0	62.0

HP 2010 Objective Data Source

25-1 a, b, c	STD Surveillance System (STDSS), CDC, NCHHSTP.
25-1 d	National Job Training Program.
25-2 a, b	STD Surveillance System (STDSS), CDC, NCHHSTP.
25-3	STD Surveillance System (STDSS), CDC, NCHHSTP.
25-4	National Health and Nutrition Examination Survey (NHANES), CDC, NCHS.
25-6	National Survey of Family Growth (NSFG), CDC, NCHS.
25-7	National Survey of Family Growth (NSFG), CDC, NCHS.
25-9	STD Surveillance System, CDC, NCHHSTP, National Vital Statistics System- Natality (NVSS-N), CDC, NCHS.
25-11 a, b, c	Youth Risk Behavior Surveillance System (YRBSS), CDC, NCCDPHP.
25-13	Survey of STD Programs, National Coalition of STD Directors (NCSDD); IHS.
25-16 a, b	Healthcare Effectiveness Data and Information Set (HEDIS), National Committee for Quality Assurance (NCQA).

*Overall positivity not adjusted for changes in laboratory test method and associated increases in test sensitivity.

[†]Data for years 1999-2004.

[‡]Data shown are for 2002, the most recent year these data were released. Beginning in 2009, data will be released every 2 years.

NOTE: Healthy People 2010 developmental objectives are not addressed in this report.

NA=Not available.

Table A4. Government Performance Results Act (GPRA) Sexually Transmitted Diseases Goals and Measures

<i>GPRA Goals</i>	<i>Baseline</i>	<i>Actual Performance</i>		<i>Long-Term Goal</i>
	<i>2002</i>	<i>2006</i>	<i>2007</i>	<i>2010</i>
Goal 1: Reduction in PID (as measured by initial visits to physicians in women 15–44-years of age)	197,000	106,000	146,000	168,000
a. Prevalence of Chlamydia in high-risk women 25 years	10.1%*	13.1%*	13.2%*	15.1%
b. Prevalence of Chlamydia in women 25 years in family planning clinics	5.6%*	6.7%*	6.9%*	7.4%
c. Incidence of Gonorrhea/100,000 population in women 15–44 years of age	279	290	290	296
Goal 2: Elimination of Syphilis (as measured by incidence of P&S Syphilis/100,000 population)	2.4	3.3	3.8	2.2
a. Incidence of P&S Syphilis/100,000 population - men	3.8	5.6	6.6	7.2
b. Incidence of P&S Syphilis/100,000 population - women	1.1	1.0	1.1	1.2
c. Incidence of Congenital Syphilis/100,000 live births	10.2	9.3	10.5	9.4
d. Black:white rate ratio of P&S Syphilis	8:1	5.9:1	7:1	6.7:1

<i>GPRA Goals</i>	<i>Data Source</i>
1	National Disease and Therapeutic Index (IMS Health).
1–a	National Job Training Program.
1–b	Regional Infertility Prevention Projects (IPP).
1–c	STD Surveillance System (STDSS), CDC, NCHHSTP.
2	STD Surveillance System (STDSS), CDC, NCHHSTP.
2–a	STD Surveillance System (STDSS), CDC, NCHHSTP.
2–b	STD Surveillance System (STDSS), CDC, NCHHSTP.
2–c	STD Surveillance System (STDSS), CDC, NCHHSTP.
2–d	STD Surveillance System (STDSS), CDC, NCHHSTP.

*Median state-specific chlamydia prevalence among women was not adjusted for changes in laboratory test method and associated increases in test sensitivity.

STD Surveillance Case Definitions

PART 1. CASE DEFINITIONS¹ FOR NATIONALLY NOTIFIABLE INFECTIOUS DISEASES

Chancroid (Revised 9/96)

Clinical description

A sexually transmitted disease characterized by painful genital ulceration and inflammatory inguinal adenopathy. The disease is caused by infection with *Haemophilus ducreyi*.

Laboratory criteria for diagnosis

- Isolation of *H. ducreyi* from a clinical specimen

Case classification

Probable: a clinically compatible case with both a) no evidence of *Treponema pallidum* infection by darkfield microscopic examination of ulcer exudate or by a serologic test for syphilis performed 7 days after onset of ulcers and b) either a clinical presentation of the ulcer(s) not typical of disease caused by herpes simplex virus (HSV) or a culture negative for HSV.

Confirmed: a clinically compatible case that is laboratory confirmed

Chlamydia trachomatis, Genital Infections (Revised 9/96)

Clinical description

Infection with *Chlamydia trachomatis* may result in urethritis, epididymitis, cervicitis, acute salpingitis, or other syndromes when sexually transmitted; however, the infection is often asymptomatic in women. Perinatal infections may result in inclusion conjunctivitis and pneumonia in newborns. Other syndromes caused by *C. trachomatis* include lymphogranuloma venereum (see Lymphogranuloma Venereum) and trachoma.

Laboratory criteria for diagnosis

- Isolation of *C. trachomatis* by culture or
- Demonstration of *C. trachomatis* in a clinical specimen by detection of antigen or nucleic acid

Case classification

Confirmed: a case that is laboratory confirmed

Gonorrhea (Revised 9/96)

Clinical description

A sexually transmitted infection commonly manifested by urethritis, cervicitis, or salpingitis. Infection may be asymptomatic.

Laboratory criteria for diagnosis

- Isolation of typical gram-negative, oxidase-positive diplococci (presumptive *Neisseria gonorrhoeae*) from a clinical specimen, or

- Demonstration of *N. gonorrhoeae* in a clinical specimen by detection of antigen or nucleic acid, or
- Observation of gram-negative intracellular diplococci in a urethral smear obtained from a male

Case classification

Probable: a) demonstration of gram-negative intracellular diplococci in an endocervical smear obtained from a female or b) a written morbidity report of gonorrhea submitted by a physician

Confirmed: a case that is laboratory confirmed

Syphilis (All Definitions Revised 9/96)

Syphilis is a complex sexually transmitted disease that has a highly variable clinical course. Classification by a clinician with expertise in syphilis may take precedence over the following case definitions developed for surveillance purposes.

Syphilis, primary

Clinical description

A stage of infection with *Treponema pallidum* characterized by one or more chancres (ulcers); chancres might differ considerably in clinical appearance.

Laboratory criteria for diagnosis

- Demonstration of *T. pallidum* in clinical specimens by darkfield microscopy, direct fluorescent antibody (DFA-TP), or equivalent methods

Case classification

Probable: a clinically compatible case with one or more ulcers (chancres) consistent with primary syphilis and a reactive serologic test (nontreponemal: Venereal Disease Research Laboratory [VDRL] or rapid plasma reagin [RPR]; treponemal: fluorescent treponemal antibody absorbed [FTA-ABS] or microhemagglutination assay for antibody to *T. pallidum* [MHA-TP])

Confirmed: a clinically compatible case that is laboratory confirmed

Syphilis, secondary

Clinical description

A stage of infection caused by *T. pallidum* and characterized by localized or diffuse mucocutaneous lesions, often with generalized lymphadenopathy. The primary chancre may still be present.

Laboratory criteria for diagnosis

- Demonstration of *T. pallidum* in clinical specimens by darkfield microscopy, DFATP, or equivalent methods

Case classification

Probable: a clinically compatible case with a nontreponemal (VDRL or RPR) titer 4

Confirmed: a clinically compatible case that is laboratory confirmed

Syphilis, latent

Clinical description

A stage of infection caused by *T. pallidum* in which organisms persist in the body of the infected person without causing symptoms or signs. Latent syphilis is subdivided into early, late, and unknown categories based on the duration of infection.

Case classification

Probable: no clinical signs or symptoms of syphilis and the presence of one of the following:

- No past diagnosis of syphilis, a reactive nontreponemal test (i.e., VDRL or RPR), and a reactive treponemal test (i.e., FTA-ABS or MHA-TP)
- A past history of syphilis therapy and a current nontreponemal test titer demonstrating fourfold or greater increase from the last nontreponemal test titer

Syphilis, early latent

Clinical description

A subcategory of latent syphilis. When initial infection has occurred within the previous 12 months, latent syphilis is classified as early latent.

Case classification

Probable: latent syphilis (see Syphilis, latent) in a person who has evidence of having acquired the infection within the previous 12 months based on one or more of the following criteria:

- Documented seroconversion or fourfold or greater increase in titer of a nontreponemal test during the previous 12 months
- A history of symptoms consistent with primary or secondary syphilis during the previous 12 months
- A history of sexual exposure to a partner who had confirmed or probable primary or secondary syphilis or probable early latent syphilis (documented independently as duration 1 year)
- Reactive nontreponemal and treponemal tests from a person whose only possible exposure occurred within the preceding 12 months

Syphilis, late latent

Clinical description

A subcategory of latent syphilis. When initial infection has occurred 1 year previously, latent syphilis is classified as late latent.

Case classification

Probable: latent syphilis (see Syphilis, latent) in a patient who has no evidence of having acquired the disease within the preceding 12 months (see Syphilis, early latent) and whose age and titer do not meet the criteria specified for latent syphilis of unknown duration.

Syphilis, latent, of unknown duration

Clinical description

A subcategory of latent syphilis. When the date of initial infection cannot be established as having occurred within the previous year and the patient's age and titer meet criteria described below, latent syphilis is classified as latent syphilis of unknown duration.

Case classification

Probable: latent syphilis (see Syphilis, latent) that does not meet the criteria for early latent syphilis, and the patient is aged 13–35 years and has a nontreponemal titer 32

Neurosyphilis

Note

Since neurosyphilis can occur at almost any stage of syphilis, between 1996 and 2005, it was classified and reported as one of several mutually exclusive stages of syphilis. In 2005, the Division of STD Prevention requested that STD control programs discontinue classifying and reporting neurosyphilis as a distinct stage of syphilis. Since 2005, if the patient has confirmed or probably neurosyphilis, the case should be reported as the appropriate state of syphilis and neurological manifestations should be noted.

Clinical description

Evidence of central nervous system infection with *T. pallidum*

Laboratory criteria for diagnosis

- A reactive serologic test for syphilis and reactive VDRL in cerebrospinal fluid (CSF)

Case classification

Probable: syphilis of any stage, a negative VDRL in CSF, and both the following:

- Elevated CSF protein or leukocyte count in the absence of other known causes of these abnormalities
- Clinical symptoms or signs consistent with neurosyphilis without other known causes for these clinical abnormalities

Confirmed: syphilis of any stage that meets the laboratory criteria for neurosyphilis

Syphilis, late, with clinical manifestations other than neurosyphilis (late benign syphilis and cardiovascular syphilis)

Clinical description

Clinical manifestations of late syphilis other than neurosyphilis may include inflammatory lesions of the cardiovascular system, skin, and bone. Rarely, other structures (e.g., the upper and lower respiratory tracts, mouth, eye, abdominal organs, reproductive organs, lymph nodes, and skeletal muscle) may be involved. Late syphilis usually becomes clinically manifest only after a period of 15–30 years of untreated infection.

Laboratory criteria for diagnosis

Demonstration of *T. pallidum* in late lesions by fluorescent antibody or special stains (although organisms are rarely visualized in late lesions)

Case classification

Probable: characteristic abnormalities or lesions of the cardiovascular system, skin, bone, or other structures with a reactive treponemal test, in the absence of other known causes of these abnormalities, and without CSF abnormalities and clinical symptoms or signs consistent with neurosyphilis

Confirmed: a clinically compatible case that is laboratory confirmed

Comment

Analysis of CSF for evidence of neurosyphilis is necessary in the evaluation of late syphilis with clinical manifestations.

Syphilitic Stillbirth

Clinical description

A fetal death that occurs after a 20-week gestation or in which the fetus weighs \leq 500 g and the mother had untreated or inadequately treated* syphilis at delivery

Comment

For reporting purposes, syphilitic stillbirths should be reported as cases of congenital syphilis.

Syphilis, Congenital (Revised 9/96)

Clinical description

A condition caused by infection in utero with *Treponema pallidum*. A wide spectrum of severity exists, and only severe cases are clinically apparent at birth. An infant or child (aged \leq 2 years) may have signs such as hepatosplenomegaly, rash, condyloma lata, snuffles, jaundice (nonviral hepatitis), pseudoparalysis, anemia, or edema (nephrotic syndrome and/or malnutrition). An older child may have stigmata (e.g., interstitial keratitis, nerve deafness, anterior bowing of shins, frontal bossing, mulberry molars, Hutchinson teeth, saddle nose, rhagades, or Clutton joints).

Laboratory criteria for diagnosis

Demonstration of *T. pallidum* by darkfield microscopy, fluorescent antibody, or other specific stains in specimens from lesions, placenta, umbilical cord, or autopsy material

Case classification

Probable: a condition affecting an infant whose mother had untreated or inadequately treated* syphilis at delivery, regardless of signs in the infant, or an infant or child who has a reactive treponemal test for syphilis and any one of the following:

- Any evidence of congenital syphilis on physical examination
- Any evidence of congenital syphilis on radiographs of long bones
- A reactive cerebrospinal fluid (CSF) venereal disease research laboratory (VDRL)
- An elevated CSF cell count or protein (without other cause)
- A reactive fluorescent treponemal antibody absorbed—19S-IgM antibody test or IgM enzyme-linked immunosorbent assay

Confirmed: a case that is laboratory confirmed

Comment

Congenital and acquired syphilis may be difficult to distinguish when a child is seropositive after infancy. Signs of congenital syphilis may not be obvious, and stigmata may not yet have developed. Abnormal values for CSF VDRL, cell count, and protein, as well as IgM antibodies, may be found in either congenital or acquired syphilis. Findings on radiographs of long bones may help because radiographic changes in the metaphysis and epiphysis are considered classic signs of congenitally acquired syphilis. The decision may ultimately be based on maternal history and clinical judgment. In a young child, the possibility of sexual abuse should be considered as a cause of acquired rather than congenital syphilis, depending on the clinical picture. For reporting purposes, congenital syphilis includes cases of congenitally acquired syphilis among infants and children as well as syphilitic stillbirths.

*Inadequate treatment consists of any nonpenicillin therapy or penicillin administered \geq 30 days before delivery.

PART 2. CASE DEFINITIONS¹ FOR NON-NOTIFIABLE INFECTIOUS DISEASES

Genital Herpes (Herpes Simplex Virus) (Revised 9/96)

Clinical description

A condition characterized by visible, painful genital or anal lesions

Laboratory criteria for diagnosis

- Isolation of herpes simplex virus from cervix, urethra, or anogenital lesion, or
- Demonstration of virus by antigen detection technique in clinical specimens from cervix, urethra, or anogenital lesion, or
- Demonstration of multinucleated giant cells on a Tzanck smear of scrapings from an anogenital lesion

Case classification

Probable: a clinically compatible case (in which primary and secondary syphilis have been excluded by appropriate serologic tests and darkfield microscopy, when available) with either a diagnosis of genital herpes based on clinical presentation (without laboratory confirmation) or a history of one or more previous episodes of similar genital lesions

Confirmed: a clinically compatible case that is laboratory confirmed

Comment

Genital herpes should be reported only once per patient. The first diagnosis for a patient with no previous diagnosis should be reported.

Genital Warts (Revised 9/96)

Clinical description

An infection characterized by the presence of visible, exophytic (raised) growths on the internal or external genitalia, perineum, or perianal region

Laboratory criteria for diagnosis

- Histopathologic changes characteristic of human papillomavirus infection in specimens obtained by biopsy or exfoliative cytology or
- Demonstration of virus by antigen or nucleic acid detection in a lesion biopsy

Case classification

Probable: a clinically compatible case without histopathologic diagnosis and without microscopic or serologic evidence that the growth is the result of secondary syphilis

Confirmed: a clinically compatible case that is laboratory confirmed

Comment

Genital warts should be reported only once per patient. The first diagnosis for a patient with no previous diagnosis should be reported.

Granuloma Inguinale

Clinical description

A slowly progressive ulcerative disease of the skin and lymphatics of the genital and perianal area caused by infection with *Calymmatobacterium granulomatis*. A clinically compatible case would have one or more painless or minimally painful granulomatous lesions in the anogenital area.

Laboratory criteria for diagnosis

- Demonstration of intracytoplasmic Donovan bodies in Wright or Giemsa-stained smears or biopsies of granulation tissue

Case classification

Confirmed: a clinically compatible case that is laboratory confirmed

Lymphogranuloma Venereum**Clinical description**

Infection with L1, L2, or, L3 serovars of *Chlamydia trachomatis* may result in a disease characterized by genital lesions, suppurative regional lymphadenopathy, or hemorrhagic proctitis. The infection is usually sexually transmitted.

Laboratory criteria for diagnosis

- Isolation of *C. trachomatis*, serotype L1, L2, or L3 from clinical specimen, or
- Demonstration by immunofluorescence of inclusion bodies in leukocytes of an inguinal lymph node (bubo) aspirate, or
- Positive microimmunofluorescent serologic test for a lymphogranuloma venereum strain of *C. trachomatis*

Case classification

Probable: a clinically compatible case with one or more tender fluctuant inguinal lymph nodes or characteristic proctogenital lesions with supportive laboratory findings of a single *C. trachomatis* complement fixation titer of ≥ 64

Confirmed: a clinically compatible case that is laboratory confirmed

Mucopurulent Cervicitis (Revised 9/96)**Clinical description**

Cervical inflammation that is not the result of infection with *Neisseria gonorrhoeae* or *Trichomonas vaginalis*. Cervical inflammation is defined by the presence of one of the following criteria:

- Mucopurulent secretion (from the endocervix) that is yellow or green when viewed on a white, cotton-tipped swab (positive swab test)
- Induced endocervical bleeding (bleeding when the first swab is placed in the endocervix)

Laboratory criteria for diagnosis

- No evidence of *N. gonorrhoeae* by culture, Gram stain, or antigen or nucleic acid detection, and no evidence of *T. vaginalis* on wet mount

Case classification

Confirmed: a clinically compatible case in a female who does not have either gonorrhea or trichomoniasis

Comment

Mucopurulent cervicitis (MPC) is a clinical diagnosis of exclusion. The syndrome may result from infection with any of several agents (see *Chlamydia trachomatis*, Genital Infections). If gonorrhea, trichomoniasis, and chlamydia are excluded, a clinically compatible illness should be classified as

MPC. An illness in a female that meets the case definition of MPC and *C. trachomatis* infection should be classified as chlamydia.

Nongonococcal Urethritis (Revised 9/96)

Clinical description

Urethral inflammation that is not the result of infection with *Neisseria gonorrhoeae*. Urethral inflammation may be diagnosed by the presence of one of the following criteria:

- A visible abnormal urethral discharge, or
- A positive leukocyte esterase test from a male aged ≥ 60 years who does not have a history of kidney disease or bladder infection, prostate enlargement, urogenital anatomic anomaly, or recent urinary tract instrumentation, or
- Microscopic evidence of urethritis (≥ 5 white blood cells per high-power field) on a Gram stain of a urethral smear

Laboratory criteria for diagnosis

- No evidence of *N. gonorrhoeae* infection by culture, Gram stain, or antigen or nucleic acid detection

Case classification

Confirmed: a clinically compatible case in a male in whom gonorrhea is not found, either by culture, Gram stain, or antigen or nucleic acid detection

Comment

Nongonococcal urethritis (NGU) is a clinical diagnosis of exclusion. The syndrome may result from infection with any of several agents (see *Chlamydia trachomatis*, Genital Infection). If gonorrhea and chlamydia are excluded, a clinically compatible illness should be classified as NGU. An illness in a male that meets the case definition of NGU and *C. trachomatis* infection should be classified as chlamydia.

Pelvic Inflammatory Disease (Revised 9/96)

Clinical case definition

A clinical syndrome resulting from the ascending spread of microorganisms from the vagina and endocervix to the endometrium, fallopian tubes, and/or contiguous structures. In a female who has lower abdominal pain and who has not been diagnosed as having an established cause other than pelvic inflammatory disease (PID) (e.g., ectopic pregnancy, acute appendicitis, and functional pain), all the following clinical criteria must be present:

- Lower abdominal tenderness, and
- Tenderness with motion of the cervix, and
- Adnexal tenderness

In addition to the preceding criteria, at least one of the following findings must also be present:

- Meets the surveillance case definition of *C. trachomatis* infection or gonorrhea
- Temperature ≥ 100.4 F (≥ 38.0 C)
- Leukocytosis $\geq 10,000$ white blood cells/mm³
- Purulent material in the peritoneal cavity obtained by culdocentesis or laparoscopy

- Pelvic abscess or inflammatory complex detected by bimanual examination or by sonography
- Patient is a sexual contact of a person known to have gonorrhea, chlamydia, or nongonococcal urethritis

Case classification

Confirmed: a case that meets the clinical case definition

Comment

For reporting purposes, a clinician's report of PID should be counted as a case.

¹ Centers for Disease Control and Prevention. Case definitions for infectious conditions under public health surveillance, 1997. *MMWR* 1997;46(No. RR-10;1).

STD Project Directors, STD Program Managers, State and Territorial Epidemiologists and Laboratory Directors

We gratefully acknowledge the contributions of state STD project directors, STD program managers, state and territorial epidemiologists and laboratory directors. The persons listed were in the positions shown as of November 4, 2008.

<u>State/City/Outlying Area</u>	<u>STD Project Directors</u>	<u>STD Program Managers</u>	<u>State Epidemiologists</u>	<u>Laboratory Directors</u>
Alabama	Charles Woernle	Sandra Langston	Charles Woernle	William Callan
Alaska	Mollie Rosier	Donna Cecere	Joe McLaughlin	Bernard Jilly
Arizona	Judy Norton	Alfonso Urquidi	Kenneth Komatsu	Victor Waddell
Arkansas	Kellye McCartney	Mark Barnes	Jim Phillips	Glen Baker
California	Gail Bolan	Romni Neiman	Gilberto Chavez	Paul Kimsey
Los Angeles	Peter Kerndt	Mary Hayes	Gilberto Chavez	Sue Sabat
San Francisco	Jeffrey Klausner	Wendy Wolf	Gilberto Chavez	Sally Liska
Colorado	Maureen Bush	Ralph Wilmoth	Ken Gershman	David Butcher
Connecticut	James Hadler	Heidi Jenkins	Matthew Cartter	John Fontana
Delaware	James Welch	Catherine Mosley	Ashley Love	Jane Getchell
District of Columbia	John Heath	John Heath	John Davies-Cole	Maurice Knuckles
Florida	Russell Eggert	Karla Schmitt	Richard Hopkins	Max Salfinger
Georgia	Vacant	Russ Cantrell	John Horan	Elizabeth Franko
Hawaii	Peter Whiticar	Roy Ohye	Sarah Park	A. Christian Whelen
Idaho	Richard Armstrong	Kathy Cohen	Christine G. Hahn	Richard Hudson
Illinois	Charlie Rabins	Ed Renier	Craig Conover (Acting)	Tom Johnson
Chicago	Will Wong	John Paffel	Craig Conover (Acting)	Susan Gerber
Indiana	Jerry Burkman	Dawne Rekas	James Howell	Judith Lovchik
Iowa	Mary Jones	Karen Thompson	Patricia Quinlisk	Christopher Atchison
Kansas	Brenda Walker	Derek Coppedge	Charles Hunt	Patrick Williams
Kentucky	Robert Brawley	Sheri White	Kraig Humbaugh	Stephanie Gibson
Louisiana	Lisa Ann Longfellow	Lisa Ann Longfellow	Raoult Ratard	Stephen Martin
Maine	Bob Woods	Jennah Godo	Kathleen Gensheimer	John Krueger
Maryland	Barbara Conrad	Glen Oltoff	David Blythe	Jack DeBoy
Baltimore	Laura Herrera	Victoria Stovall	David Blythe	Jack DeBoy
Massachusetts	Thomas Bertrand	Thomas Bertrand	Alfred DeMaria	Mary Gilchrist
Michigan	Mark Miller	Mark Miller	Corrine Miller	Frances Downes
Minnesota	Julia Ashley	Julia Ashley	Ruth Lynfield	Joanne Bartkus
Mississippi	Craig Thompson	Craig Thompson	Mary Currier	K. Mills McNeill
Missouri	Michael Herbert	Michael Herbert	Sarah Patrick	Eric Blank
Montana	Laurie Kops	Lisa Underwood	Steve Helgerson	Anne Weber
Nebraska	Phil Medina	Phil Medina	Thomas Safranek	Steve Hinrichs
Nevada	Richard Whitley	Julia Spaulding	Ishan Azzam	L. Dee Brown
New Hampshire	Denise Rondeau	Drew Thomits	Jose Montero	Christine Bean
New Jersey	Janet DeGraaf	Patricia Masou	Christina Tan	Dennis Flynn
New Mexico	Daryl Smith	Daryl Smith	C. Mack Sewell	David Mills
New York	F. Bruce Coles	Dennis Murphy	Perry Smith	Lawrence Sturman
New York City	Susan Blank	Steve Rubin	Perry Smith	Sara Beatrice
North Carolina	Evelyn Foust	Evelyn Foust	Jeffrey Engel	Leslie Wolf
North Dakota	Kimberly Weis	Julie Goplin	Kirby Kruger	Myra Kosse
Ohio	Bill Tiedemann	Amy LaGessee	Forrest Smith	Tammy Bannerman
Oklahoma	Michael Harmon	Chang Lee	Kristy Bradley	Garry McKee
Oregon	Vada Latin	Doug Harger	Melvin Kohn	Michael Skeels
Pennsylvania	Weston Ruhrig	Steve Kowalewski	Veronica Urdaneta	S.I. Shahied
Philadelphia	Caroline Johnson	Melinda Salmon	Veronica Urdaneta	Kerry Buchs
Rhode Island	Utpala Bandy	Michael Gosciminski	Utpala Bandy	Ewa King
South Carolina	Dorothy Waln (Acting)	Vickie Boazman-Holmes (Acting)	James Gibson	Arthur Wozniak
South Dakota	Laurie Gill	David Morgan	Lon Kightlinger	Michael Smith
Tennessee	Jeanee Seals	Jane Russell	Tim Jones	David Smalley
Texas	Felipe Rocha	Jim Lee	Vincent Fonseca	Susan Neill
Utah	Jennifer Brown	Tim Lane	Robert Rolfs	Patrick Luedtke
Vermont	Daniel Daltry	Daniel Daltry	Patsy Kelso	Mary Celotti
Virginia	Kathryn Hafford	Kathryn Hafford (Acting)	Carl Armstrong	James Pearson
Washington	Mark Aubin	Mark Aubin	Marcia Goldoft	Romesch Gautom
West Virginia	Caroline Williams	Caroline Williams	Loretta Haddy	Andrea Labik
Wisconsin	Sandra Breitborde	Anthony Wade	Jeffrey Davis	Charles Brokopp
Wyoming	Brownen Anderson	Canyon Hardesty	Tracy Murphy	Richard Harris
American Samoa	Utoofili Maga	Sarona Maae	Sharmain Edwards	Utoofili Mago
Federated States of Micronesia	Vita Skilling	Johnny Hebel	Nena Nena	
Government of the Marshall Islands	Justina Langidrik	Zachraias Zachraias		Paul Lalita
Northern Marianas (CNMI)	Joseph Villagomez	John Moreno	James Hosfschneider	Stephanie Hart-Gouleau
Guam	Josie O'Mallan	Bernadette Schumann	Robert Haddock	Josie O'Mallan
Puerto Rico	Greduvel Duran-Guzman	Trinidad Garcia	Enid Garcia	Francisco Toro
Republic of Palau	Victor Yano	Johana Ngiruchelbad	Julie Erb-Alvarez	Francis Permeteet
Virgin Islands	Darlene Carty	Taetia Phillips-Dorsett	Darlene Carty	Joseph Mark