

Alabama, 2007

Primary and secondary syphilis

- **CASES:** Alabama reported 380 cases in 2007.
- **RATES:** Alabama ranked 3 among 50 states, DC, and 3 territories with 8.3 cases per 100,000 population compared to the total rate of 3.8 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Alabama, the rate among males was 10.6 per 100,000 population compared to the total male rate of 6.6 per 100,000.
- The rate among females was 6.0 per 100,000 compared to the total female rate of 1.1 per 100,000.

Rates of primary and secondary syphilis, by race/ethnicity


In Alabama, the race/ethnicity adjusted rates per 100,000 population were 2.7 among whites, 23.2 among blacks, 9.7 among Hispanics, 2.2 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.

- The rate among blacks was 8.6 times that of whites.
- The rate among Hispanics was 3.6 times that of whites.

Congenital syphilis

- **CASES:** Alabama reported 9 cases in 2007.
- **RATES:** Alabama ranked 11 among 31 states, DC, and 2 territories reporting congenital syphilis cases with a rate of 15.1 cases per 100,000 live births compared to the total rate of 10.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Alabama, 1998-2007



Total P&S Syphilis Rates by County, Alabama, 2007

