

District of Columbia, 2007

Primary and secondary syphilis

- **CASES:** District of Columbia reported 178 cases in 2007.
- **RATES:** District of Columbia ranked 1 among 50 states, DC, and 3 territories with 30.6 cases per 100,000 population compared to the total rate of 3.8 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In District of Columbia, the rate among males was 63.4 per 100,000 population compared to the total male rate of 6.6 per 100,000.
- The rate among females was 1.6 per 100,000 compared to the total female rate of 1.1 per 100,000.

Rates of primary and secondary syphilis, by race/ethnicity


In District of Columbia, the race/ethnicity adjusted rates per 100,000 population were 28.5 among whites, 31.6 among blacks, 39.8 among Hispanics, 15.1 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.

- The rate among blacks was 1.1 times that of whites.
- The rate among Hispanics was 1.4 times that of whites.

Congenital syphilis

- **CASES:** District of Columbia reported 1 case in 2007.
- **RATES:** District of Columbia ranked 12 among 31 states, DC, and 2 territories reporting congenital syphilis cases with a rate of 12.6 cases per 100,000 live births compared to the total rate of 10.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, District of Columbia, 1998-2007



Total P&S Syphilis Rate, District of Columbia, 2007

