

# Alabama, 2008

## Primary and secondary syphilis

- CASES: Alabama reported 449 cases in 2008.
- RATES: Alabama ranked 3 among 50 states; Washington, DC; and 3 territories with 9.7 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Alabama, the rate among males was 12.4 per 100,000 population compared to the U.S. male rate of 7.6 per 100,000.
- The rate among females was 7.2 per 100,000 compared to the U.S. female rate of 1.5 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Alabama, the race/ethnicity adjusted rates per 100,000 population were 2.5 among whites, 28.6 among blacks, 14.4 among Hispanics, 2.1 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 11.4 times that of whites.
- The rate among Hispanics was 5.8 times that of whites.


## Congenital syphilis

- CASES: Alabama reported 12 cases in 2008.
- RATES: Alabama ranked 7 among 26 states; Washington, DC; and 1 territory reporting congenital syphilis cases with a rate of 19.0 cases per 100,000 live births compared to the U.S. rate of 10.1 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Alabama, 1999-2008



## Total P&S Syphilis Rates by County, Alabama, 2008



Rate (per 100,000 population)

