

Connecticut, 2008

Primary and secondary syphilis

- CASES: Connecticut reported 34 cases in 2008.
- RATES: Connecticut ranked 41 among 50 states; Washington, DC; and 3 territories with 1 case per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Connecticut, the rate among males was 2.0 per 100,000 population compared to the U.S. male rate of 7.6 per 100,000.
- The rate among females was 0.0 per 100,000 compared to the U.S. female rate of 1.5 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Connecticut, the race/ethnicity adjusted rates per 100,000 population were 0.7 among whites, 2.6 among blacks, 1.7 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 3.7 times that of whites.
- The rate among Hispanics was 2.4 times that of whites.

Congenital syphilis

- CASES: Connecticut reported 2 cases in 2008.
- RATES: Connecticut ranked 20 among 26 states; Washington, DC; and 1 territory reporting congenital syphilis cases with a rate of 4.8 cases per 100,000 live births compared to the U.S. rate of 10.1 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Connecticut, 1999-2008



Total P&S Syphilis Rates by County, Connecticut, 2008

