

New York, 2008

Primary and secondary syphilis

- CASES: New York reported 1,217 cases in 2008.
- RATES: New York ranked 8 among 50 states; Washington, DC; and 3 territories with 6.3 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In New York, the rate among males was 12.4 per 100,000 population compared to the U.S. male rate of 7.6 per 100,000.
- The rate among females was 0.5 per 100,000 compared to the U.S. female rate of 1.5 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In New York, the race/ethnicity adjusted rates per 100,000 population were 3.0 among whites, 17.9 among blacks, 9.6 among Hispanics, 1.8 among Asian/Pacific Islanders, and 6.9 among American Indians/Alaska Natives.
- The rate among blacks was 6 times that of whites.
- The rate among Hispanics was 3.2 times that of whites.
- The rate among American Indians/Alaska Natives was 2.3 times that of whites.

Congenital syphilis

- CASES: New York reported 23 cases in 2008.
- RATES: New York ranked 13 among 26 states; Washington, DC; and 1 territory reporting congenital syphilis cases with a rate of 9.2 cases per 100,000 live births compared to the U.S. rate of 10.1 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, New York, 1999-2008



Total P&S Syphilis Rates by County, New York, 2008

