

# Ohio, 2008

## Primary and secondary syphilis

- CASES: Ohio reported 351 cases in 2008.
- RATES: Ohio ranked 21 among 50 states; Washington, DC; and 3 territories with 3.1 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Ohio, the rate among males was 5.2 per 100,000 population compared to the U.S. male rate of 7.6 per 100,000.
- The rate among females was 1.1 per 100,000 compared to the U.S. female rate of 1.5 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Ohio, the race/ethnicity adjusted rates per 100,000 population were 1.7 among whites, 12.3 among blacks, 3.5 among Hispanics, 1.6 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 7.2 times that of whites.
- The rate among Hispanics was 2.1 times that of whites.


## Congenital syphilis

- CASES: Ohio reported 3 cases in 2008.
- RATES: Ohio ranked 27 among 26 states; Washington, DC; and 1 territory reporting congenital syphilis cases with a rate of 2.0 cases per 100,000 live births compared to the U.S. rate of 10.1 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Ohio, 1999-2008



## Total P&S Syphilis Rates by County, Ohio, 2008



Rate (per 100,000 population)

