

Oklahoma, 2008

Primary and secondary syphilis

- CASES: Oklahoma reported 86 cases in 2008.
- RATES: Oklahoma ranked 26 among 50 states; Washington, DC; and 3 territories with 2.4 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Oklahoma, the rate among males was 3.4 per 100,000 population compared to the U.S. male rate of 7.6 per 100,000.
- The rate among females was 1.4 per 100,000 compared to the U.S. female rate of 1.5 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Oklahoma, the race/ethnicity adjusted rates per 100,000 population were 0.9 among whites, 13.6 among blacks, 4.6 among Hispanics, 1.5 among Asian/Pacific Islanders, and 2.6 among American Indians/Alaska Natives.
- The rate among blacks was 15.1 times that of whites.
- The rate among Hispanics was 5.1 times that of whites.
- The rate among Asian/Pacific Islanders was 1.7 times that of whites.
- The rate among American Indians/Alaska Natives was 2.9 times that of whites.


Congenital syphilis

- CASES: Oklahoma reported 3 cases in 2008.
- RATES: Oklahoma ranked 18 among 26 states; Washington, DC; and 1 territory reporting congenital syphilis cases with a rate of 5.6 cases per 100,000 live births compared to the U.S. rate of 10.1 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Oklahoma, 1999-2008



Total P&S Syphilis Rates by County, Oklahoma, 2008



Rate (per 100,000 population)

