Hawaii, 2009

Primary and secondary syphilis

- CASES: Hawaii reported 33 cases in 2009.
- RATES: Hawaii ranked 27 among 50 states; Washington, DC; and 3 territories with 2.6 cases per 100,000 population compared to the U.S. rate of 4.6 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Hawaii, the rate among males was 4.2 per 100,000 population compared to the U.S. male rate of 7.8 per 100,000.
- The rate among females was 0.9 per 100,000 compared to the U.S. female rate of 1.4 per 100,000.

Rates of primary and secondary syphilis, by race/ethnicity

- In Hawaii, the race/ethnicity adjusted rates per 100,000 population were 4.7 among whites, 7.2 among blacks, 3.6 among Hispanics, 1.3 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 1.5 times that of whites.

Congenital syphilis

- CASES: Hawaii reported 1 case in 2009.
- RATES: Hawaii ranked 23 among among 34 areas (33 states and 1 territory) reporting congenital syphilis cases with a rate of 5.3 cases per 100,000 live births compared to the U.S. rate of 10.0 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Hawaii, 2000-2009



Total P&S Syphilis Rates by County, Hawaii, 2009