

Ohio, 2009

Primary and secondary syphilis

- CASES: Ohio reported 360 cases in 2009.
- RATES: Ohio ranked 20 among 50 states; Washington, DC; and 3 territories with 3.1 cases per 100,000 population compared to the U.S. rate of 4.6 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Ohio, the rate among males was 5.3 per 100,000 population compared to the U.S. male rate of 7.8 per 100,000.
- The rate among females was 1.1 per 100,000 compared to the U.S. female rate of 1.4 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Ohio, the race/ethnicity adjusted rates per 100,000 population were 1.4 among whites, 15.4 among blacks, 2.0 among Hispanics, 1.5 among Asian/Pacific Islanders, and 3.3 among American Indians/Alaska Natives.
- The rate among blacks was 11 times that of whites.
- The rate among Hispanics was 1.4 times that of whites.
- The rate among Asian/Pacific Islanders was 1.1 times that of whites.
- The rate among American Indians/Alaska Natives was 2.4 times that of whites.

Congenital syphilis

- CASES: Ohio reported 7 cases in 2009.
- RATES: Ohio ranked 25 among among 34 areas (33 states and 1 territory) reporting congenital syphilis cases with a rate of 4.6 cases per 100,000 live births compared to the U.S. rate of 10.0 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Ohio, 2000-2009



Total P&S Syphilis Rates by County, Ohio, 2009

