

Wyoming, 2009

Primary and secondary syphilis

- CASES: Wyoming reported 3 cases in 2009.
- RATES: Wyoming ranked 44 among 50 states; Washington, DC; and 3 territories with 0.6 cases per 100,000 population compared to the U.S. rate of 4.6 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Wyoming, the rate among males was 0.7 per 100,000 population compared to the U.S. male rate of 7.8 per 100,000.
- The rate among females was 0.4 per 100,000 compared to the U.S. female rate of 1.4 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Wyoming, the race/ethnicity adjusted rates per 100,000 population were 0.6 among whites, 0.0 among blacks, 0.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.


Congenital syphilis

- CASES: Wyoming reported 0 cases in 2009.
- RATES: Wyoming had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 10.0 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Wyoming, 2000-2009



Total P&S Syphilis Rates by County, Wyoming, 2009



Rate (per 100,000 population)

