

California, 2010

Primary and secondary syphilis

- CASES: California reported 2,065 cases in 2010.
- RATES: California ranked 11 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 5.6 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In California, the rate among males was 10.8 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.4 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In California, the race/ethnicity adjusted rates per 100,000 population were 5.7 among whites, 12.8 among blacks, 5.0 among Hispanics, 1.9 among Asian/Pacific Islanders, and 2.5 among American Indians/Alaska Natives.
- The rate among blacks was 2.2 times that of whites.


Congenital syphilis

- CASES: California reported 38 cases in 2010.
- RATES: California ranked 16 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 6.7 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, California, 2001-2010



Total P&S Syphilis Rates by County, California, 2010



Rate (per 100,000 population)

