

# Alabama, 2010

## Primary and secondary syphilis

- CASES: Alabama reported 260 cases in 2010.
- RATES: Alabama ranked 12 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 5.5 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Alabama, the rate among males was 8.1 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 3.1 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Alabama, the race/ethnicity adjusted rates per 100,000 population were 1.1 among whites, 17.6 among blacks, 2.6 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 16 times that of whites.
- The rate among Hispanics was 2.4 times that of whites.


## Congenital syphilis

- CASES: Alabama reported 9 cases in 2010.
- RATES: Alabama ranked 9 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 13.9 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Alabama, 2001-2010



## Total P&S Syphilis Rates by County, Alabama, 2010



Rate (per 100,000 population)



# Alaska, 2010

## Primary and secondary syphilis

- CASES: Alaska reported 3 cases in 2010.
- RATES: Alaska ranked 49 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 0.4 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Alaska, the rate among males was 0.8 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.0 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Alaska, the race/ethnicity adjusted rates per 100,000 population were 0.6 among whites, 0.0 among blacks, 0.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.


## Congenital syphilis

- CASES: Alaska reported 0 cases in 2010.
- RATES: Alaska had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Alaska, 2001-2010



## Total P&S Syphilis Rates by County, Alaska, 2010





# Arkansas, 2010

## Primary and secondary syphilis

- CASES: Arkansas reported 205 cases in 2010.
- RATES: Arkansas ranked 5 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 7.1 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Arkansas, the rate among males was 8.7 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 5.6 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Arkansas, the race/ethnicity adjusted rates per 100,000 population were 2.4 among whites, 32.1 among blacks, 2.3 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 13.4 times that of whites.


## Congenital syphilis

- CASES: Arkansas reported 11 cases in 2010.
- RATES: Arkansas ranked 3 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 26.6 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Arkansas, 2001-2010



## Total P&S Syphilis Rates by County, Arkansas, 2010



# California, 2010

## Primary and secondary syphilis

- CASES: California reported 2,065 cases in 2010.
- RATES: California ranked 11 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 5.6 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In California, the rate among males was 10.8 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.4 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In California, the race/ethnicity adjusted rates per 100,000 population were 5.7 among whites, 12.8 among blacks, 5.0 among Hispanics, 1.9 among Asian/Pacific Islanders, and 2.5 among American Indians/Alaska Natives.
- The rate among blacks was 2.2 times that of whites.


## Congenital syphilis

- CASES: California reported 38 cases in 2010.
- RATES: California ranked 16 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 6.7 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, California, 2001-2010



## Total P&S Syphilis Rates by County, California, 2010



Rate (per 100,000 population)



# Colorado, 2010

## Primary and secondary syphilis

- **CASES:** Colorado reported 138 cases in 2010.
- **RATES:** Colorado ranked 29 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.7 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Colorado, the rate among males was 5.4 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.1 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Colorado, the race/ethnicity adjusted rates per 100,000 population were 2.6 among whites, 6.9 among blacks, 2.6 among Hispanics, 0.7 among Asian/Pacific Islanders, and 2.3 among American Indians/Alaska Natives.
- The rate among blacks was 2.7 times that of whites.


## Congenital syphilis

- **CASES:** Colorado reported 0 cases in 2010.
- **RATES:** Colorado had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Colorado, 2001-2010



## Total P&S Syphilis Rates by County, Colorado, 2010



Rate (per 100,000 population)



# Connecticut, 2010

## Primary and secondary syphilis

- CASES: Connecticut reported 98 cases in 2010.
- RATES: Connecticut ranked 28 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.8 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Connecticut, the rate among males was 5.4 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.3 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Connecticut, the race/ethnicity adjusted rates per 100,000 population were 2.1 among whites, 6.9 among blacks, 4.6 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 3.3 times that of whites.
- The rate among Hispanics was 2.2 times that of whites.


## Congenital syphilis

- CASES: Connecticut reported 2 cases in 2010.
- RATES: Connecticut ranked 19 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 4.8 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Connecticut, 2001-2010



## Total P&S Syphilis Rates by County, Connecticut, 2010



# Delaware, 2010

## Primary and secondary syphilis

- CASES: Delaware reported 9 cases in 2010.
- RATES: Delaware ranked 40 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 1 case per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Delaware, the rate among males was 1.9 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.2 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Delaware, the race/ethnicity adjusted rates per 100,000 population were 0.3 among whites, 3.7 among blacks, 0.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 12.3 times that of whites.


## Congenital syphilis

- CASES: Delaware reported 2 cases in 2010.
- RATES: Delaware ranked 6 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 16.4 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Delaware, 2001-2010



## Total P&S Syphilis Rates by County, Delaware, 2010



Rate (per 100,000 population)



# District of Columbia, 2010

## Primary and secondary syphilis

- **CASES:** District of Columbia reported 134 cases in 2010.
- **RATES:** District of Columbia ranked 1 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 22.3 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In District of Columbia, the rate among males was 46.6 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.6 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In District of Columbia, the race/ethnicity adjusted rates per 100,000 population were 22.1 among whites, 22.1 among blacks, 20.7 among Hispanics, 19.5 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.


## Congenital syphilis

- **CASES:** District of Columbia reported 1 case in 2010.
- **RATES:** District of Columbia ranked 13 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 11.3 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, District of Columbia, 2001-2010



## Total P&S Syphilis Rate, District of Columbia, 2010



# Florida, 2010

## Primary and secondary syphilis

- CASES: Florida reported 1,184 cases in 2010.
- RATES: Florida ranked 7 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 6.4 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Florida, the rate among males was 11.4 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 1.6 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Florida, the race/ethnicity adjusted rates per 100,000 population were 3.5 among whites, 15.9 among blacks, 7.1 among Hispanics, 0.6 among Asian/Pacific Islanders, and 3.3 among American Indians/Alaska Natives.
- The rate among blacks was 4.5 times that of whites.
- The rate among Hispanics was 2 times that of whites.


## Congenital syphilis

- CASES: Florida reported 19 cases in 2010.
- RATES: Florida ranked 14 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 7.9 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Florida, 2001-2010



## Total P&S Syphilis Rates by County, Florida, 2010



Rate (per 100,000 population)



# Georgia, 2010

## Primary and secondary syphilis

- CASES: Georgia reported 795 cases in 2010.
- RATES: Georgia ranked 3 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 8.1 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Georgia, the rate among males was 14.7 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 1.6 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Georgia, the race/ethnicity adjusted rates per 100,000 population were 2.0 among whites, 21.6 among blacks, 3.2 among Hispanics, 1.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 10.8 times that of whites.
- The rate among Hispanics was 1.6 times that of whites.


## Congenital syphilis

- CASES: Georgia reported 18 cases in 2010.
- RATES: Georgia ranked 12 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 11.9 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Georgia, 2001-2010



## Total P&S Syphilis Rates by County, Georgia, 2010



Rate (per 100,000 population)



# Hawaii, 2010

## Primary and secondary syphilis

- CASES: Hawaii reported 35 cases in 2010.
- RATES: Hawaii ranked 31 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.7 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Hawaii, the rate among males was 4.3 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 1.1 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Hawaii, the race/ethnicity adjusted rates per 100,000 population were 3.6 among whites, 4.6 among blacks, 4.3 among Hispanics, 1.6 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 1.3 times that of whites.
- The rate among Hispanics was 1.2 times that of whites.


## Congenital syphilis

- CASES: Hawaii reported 0 cases in 2010.
- RATES: Hawaii had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Hawaii, 2001-2010



## Total P&S Syphilis Rates by County, Hawaii, 2010



Rate (per 100,000 population)



# Idaho, 2010

## Primary and secondary syphilis

- CASES: Idaho reported 6 cases in 2010.
- RATES: Idaho ranked 50 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 0.4 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Idaho, the rate among males was 0.8 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.0 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Idaho, the race/ethnicity adjusted rates per 100,000 population were 0.2 among whites, 0.0 among blacks, 1.8 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among Hispanics was 9 times that of whites.

## Congenital syphilis

- CASES: Idaho reported 1 case in 2010.
- RATES: Idaho ranked 21 among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 4.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Idaho, 2001-2010



## Total P&S Syphilis Rates by County, Idaho, 2010



Rate (per 100,000 population)



# Illinois, 2010

## Primary and secondary syphilis

- **CASES:** Illinois reported 908 cases in 2010.
- **RATES:** Illinois ranked 6 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 7 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Illinois, the rate among males was 12.6 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 1.6 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Illinois, the race/ethnicity adjusted rates per 100,000 population were 2.6 among whites, 27.7 among blacks, 5.8 among Hispanics, 1.7 among Asian/Pacific Islanders, and 14.1 among American Indians/Alaska Natives.
- The rate among blacks was 10.7 times that of whites.
- The rate among Hispanics was 2.2 times that of whites.
- The rate among American Indians/Alaska Natives was 5.4 times that of whites.


## Congenital syphilis

- **CASES:** Illinois reported 27 cases in 2010.
- **RATES:** Illinois ranked 7 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 14.9 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Illinois, 2001-2010



## Total P&S Syphilis Rates by County, Illinois, 2010



Rate (per 100,000 population)



# Indiana, 2010

## Primary and secondary syphilis

- CASES: Indiana reported 175 cases in 2010.
- RATES: Indiana ranked 30 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.7 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Indiana, the rate among males was 4.9 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.6 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Indiana, the race/ethnicity adjusted rates per 100,000 population were 1.3 among whites, 16.0 among blacks, 2.9 among Hispanics, 1.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 12.3 times that of whites.
- The rate among Hispanics was 2.2 times that of whites.


## Congenital syphilis

- CASES: Indiana reported 0 cases in 2010.
- RATES: Indiana had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Indiana, 2001-2010



## Total P&S Syphilis Rates by County, Indiana, 2010



Rate (per 100,000 population)



# Iowa, 2010

## Primary and secondary syphilis

- CASES: Iowa reported 19 cases in 2010.
- RATES: Iowa ranked 45 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 0.6 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Iowa, the rate among males was 1.1 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.2 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Iowa, the race/ethnicity adjusted rates per 100,000 population were 0.6 among whites, 2.2 among blacks, 0.7 among Hispanics, 1.9 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 3.7 times that of whites.
- The rate among Hispanics was 1.2 times that of whites.
- The rate among Asian/Pacific Islanders was 3.2 times that of whites.


## Congenital syphilis

- CASES: Iowa reported 0 cases in 2010.
- RATES: Iowa had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Iowa, 2001-2010



## Total P&S Syphilis Rates by County, Iowa, 2010



Rate (per 100,000 population)



# Kansas, 2010

## Primary and secondary syphilis

- CASES: Kansas reported 19 cases in 2010.
- RATES: Kansas ranked 42 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 0.7 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Kansas, the rate among males was 1.3 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.1 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Kansas, the race/ethnicity adjusted rates per 100,000 population were 0.5 among whites, 2.8 among blacks, 0.8 among Hispanics, 0.0 among Asian/Pacific Islanders, and 3.6 among American Indians/Alaska Natives.
- The rate among blacks was 5.6 times that of whites.
- The rate among Hispanics was 1.6 times that of whites.
- The rate among American Indians/Alaska Natives was 7.2 times that of whites.


## Congenital syphilis

- CASES: Kansas reported 0 cases in 2010.
- RATES: Kansas had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Kansas, 2001-2010



## Total P&S Syphilis Rates by County, Kansas, 2010



Rate (per 100,000 population)



# Kentucky, 2010

## Primary and secondary syphilis

- **CASES:** Kentucky reported 139 cases in 2010.
- **RATES:** Kentucky ranked 24 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 3.2 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Kentucky, the rate among males was 6.2 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.4 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Kentucky, the race/ethnicity adjusted rates per 100,000 population were 2.2 among whites, 12.7 among blacks, 6.9 among Hispanics, 2.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 5.8 times that of whites.
- The rate among Hispanics was 3.1 times that of whites.


## Congenital syphilis

- **CASES:** Kentucky reported 0 cases in 2010.
- **RATES:** Kentucky had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Kentucky, 2001-2010



## Total P&S Syphilis Rates by County, Kentucky, 2010



Rate (per 100,000 population)



# Louisiana, 2010

## Primary and secondary syphilis

- CASES: Louisiana reported 546 cases in 2010.
- RATES: Louisiana ranked 2 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 12.2 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Louisiana, the rate among males was 13.0 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 10.9 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Louisiana, the race/ethnicity adjusted rates per 100,000 population were 1.7 among whites, 30.8 among blacks, 2.5 among Hispanics, 2.8 among Asian/Pacific Islanders, and 7.1 among American Indians/Alaska Natives.
- The rate among blacks was 18.1 times that of whites.
- The rate among Hispanics was 1.5 times that of whites.
- The rate among Asian/Pacific Islanders was 1.6 times that of whites.
- The rate among American Indians/Alaska Natives was 4.2 times that of whites.


## Congenital syphilis

- CASES: Louisiana reported 33 cases in 2010.
- RATES: Louisiana ranked 1 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 49.8 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Louisiana, 2001-2010



## Total P&S Syphilis Rates by County, Louisiana, 2010



Rate (per 100,000 population)



# Maine, 2010

## Primary and secondary syphilis

- CASES: Maine reported 32 cases in 2010.
- RATES: Maine ranked 35 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.4 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Maine, the rate among males was 5.0 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.0 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Maine, the race/ethnicity adjusted rates per 100,000 population were 2.4 among whites, 0.0 among blacks, 0.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.


## Congenital syphilis

- CASES: Maine reported 0 cases in 2010.
- RATES: Maine had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Maine, 2001-2010



## Total P&S Syphilis Rates by County, Maine, 2010



# Maryland, 2010

## Primary and secondary syphilis

- CASES: Maryland reported 328 cases in 2010.
- RATES: Maryland ranked 8 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 5.8 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Maryland, the rate among males was 10.9 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.9 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Maryland, the race/ethnicity adjusted rates per 100,000 population were 1.1 among whites, 15.9 among blacks, 2.7 among Hispanics, 1.6 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 14.5 times that of whites.
- The rate among Hispanics was 2.5 times that of whites.
- The rate among Asian/Pacific Islanders was 1.5 times that of whites.


## Congenital syphilis

- CASES: Maryland reported 22 cases in 2010.
- RATES: Maryland ranked 2 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 28.2 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Maryland, 2001-2010



## Total P&S Syphilis Rates by County, Maryland, 2010



# Massachusetts, 2010

## Primary and secondary syphilis

- CASES: Massachusetts reported 285 cases in 2010.
- RATES: Massachusetts ranked 17 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 4.3 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Massachusetts, the rate among males was 8.4 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.5 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Massachusetts, the race/ethnicity adjusted rates per 100,000 population were 3.2 among whites, 11.0 among blacks, 8.2 among Hispanics, 1.7 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 3.4 times that of whites.
- The rate among Hispanics was 2.6 times that of whites.


## Congenital syphilis

- CASES: Massachusetts reported 1 case in 2010.
- RATES: Massachusetts ranked 28 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 1.3 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Massachusetts, 2001-2010



## Total P&S Syphilis Rates by County, Massachusetts, 2010



Rate (per 100,000 population)



# Michigan, 2010

## Primary and secondary syphilis

- CASES: Michigan reported 235 cases in 2010.
- RATES: Michigan ranked 36 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.4 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Michigan, the rate among males was 4.3 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.5 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Michigan, the race/ethnicity adjusted rates per 100,000 population were 1.1 among whites, 9.2 among blacks, 2.4 among Hispanics, 0.8 among Asian/Pacific Islanders, and 1.5 among American Indians/Alaska Natives.
- The rate among blacks was 8.4 times that of whites.
- The rate among Hispanics was 2.2 times that of whites.
- The rate among American Indians/Alaska Natives was 1.4 times that of whites.


## Congenital syphilis

- CASES: Michigan reported 2 cases in 2010.
- RATES: Michigan ranked 26 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 1.6 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Michigan, 2001-2010



## Total P&S Syphilis Rates by County, Michigan, 2010



Rate (per 100,000 population)



# Minnesota, 2010

## Primary and secondary syphilis

- **CASES:** Minnesota reported 149 cases in 2010.
- **RATES:** Minnesota ranked 26 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.8 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Minnesota, the rate among males was 5.3 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.3 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Minnesota, the race/ethnicity adjusted rates per 100,000 population were 2.0 among whites, 15.0 among blacks, 5.3 among Hispanics, 0.5 among Asian/Pacific Islanders, and 1.6 among American Indians/Alaska Natives.
- The rate among blacks was 7.5 times that of whites.
- The rate among Hispanics was 2.7 times that of whites.


## Congenital syphilis

- **CASES:** Minnesota reported 0 cases in 2010.
- **RATES:** Minnesota had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Minnesota, 2001-2010



## Total P&S Syphilis Rates by County, Minnesota, 2010



Rate (per 100,000 population)



# Mississippi, 2010

## Primary and secondary syphilis

- CASES: Mississippi reported 228 cases in 2010.
- RATES: Mississippi ranked 4 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 7.7 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Mississippi, the rate among males was 11.1 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 4.5 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Mississippi, the race/ethnicity adjusted rates per 100,000 population were 2.0 among whites, 17.4 among blacks, 1.3 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 8.7 times that of whites.



## Congenital syphilis

- CASES: Mississippi reported 9 cases in 2010.
- RATES: Mississippi ranked 5 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 19.4 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Mississippi, 2001-2010



## Total P&S Syphilis Rates by County, Mississippi, 2010



# Missouri, 2010

## Primary and secondary syphilis

- CASES: Missouri reported 152 cases in 2010.
- RATES: Missouri ranked 33 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.5 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Missouri, the rate among males was 5.1 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.1 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Missouri, the race/ethnicity adjusted rates per 100,000 population were 1.2 among whites, 11.4 among blacks, 2.9 among Hispanics, 2.0 among Asian/Pacific Islanders, and 3.1 among American Indians/Alaska Natives.
- The rate among blacks was 9.5 times that of whites.
- The rate among Hispanics was 2.4 times that of whites.
- The rate among Asian/Pacific Islanders was 1.7 times that of whites.
- The rate among American Indians/Alaska Natives was 2.6 times that of whites.


## Congenital syphilis

- CASES: Missouri reported 2 cases in 2010.
- RATES: Missouri ranked 23 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 2.4 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Missouri, 2001-2010



## Total P&S Syphilis Rates by County, Missouri, 2010



# Montana, 2010

## Primary and secondary syphilis

- **CASES:** Montana reported 3 cases in 2010.
- **RATES:** Montana ranked 52 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 0.3 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Montana, the rate among males was 0.6 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.0 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Montana, the race/ethnicity adjusted rates per 100,000 population were 0.3 among whites, 0.0 among blacks, 0.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.


## Congenital syphilis

- **CASES:** Montana reported 0 cases in 2010.
- **RATES:** Montana had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Montana, 2001-2010



## Total P&S Syphilis Rates by County, Montana, 2010



Rate (per 100,000 population)



# Nebraska, 2010

## Primary and secondary syphilis

- **CASES:** Nebraska reported 12 cases in 2010.
- **RATES:** Nebraska ranked 43 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 0.7 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Nebraska, the rate among males was 1.0 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.3 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Nebraska, the race/ethnicity adjusted rates per 100,000 population were 0.5 among whites, 0.0 among blacks, 1.3 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among Hispanics was 2.6 times that of whites.


## Congenital syphilis

- **CASES:** Nebraska reported 0 cases in 2010.
- **RATES:** Nebraska had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Nebraska, 2001-2010



## Total P&S Syphilis Rates by County, Nebraska, 2010



Rate (per 100,000 population)



# Nevada, 2010

## Primary and secondary syphilis

- CASES: Nevada reported 130 cases in 2010.
- RATES: Nevada ranked 14 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 4.9 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Nevada, the rate among males was 9.1 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.5 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Nevada, the race/ethnicity adjusted rates per 100,000 population were 3.1 among whites, 18.4 among blacks, 4.1 among Hispanics, 1.0 among Asian/Pacific Islanders, and 3.3 among American Indians/Alaska Natives.
- The rate among blacks was 5.9 times that of whites.
- The rate among Hispanics was 1.3 times that of whites.
- The rate among American Indians/Alaska Natives was 1.1 times that of whites.



## Congenital syphilis

- CASES: Nevada reported 5 cases in 2010.
- RATES: Nevada ranked 11 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 12.1 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Nevada, 2001-2010



## Total P&S Syphilis Rates by County, Nevada, 2010



# New Hampshire, 2010

## Primary and secondary syphilis

- CASES: New Hampshire reported 22 cases in 2010.
- RATES: New Hampshire ranked 39 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 1.7 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In New Hampshire, the rate among males was 3.2 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.1 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In New Hampshire, the race/ethnicity adjusted rates per 100,000 population were 1.5 among whites, 5.4 among blacks, 2.7 among Hispanics, 7.3 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 3.6 times that of whites.
- The rate among Hispanics was 1.8 times that of whites.
- The rate among Asian/Pacific Islanders was 4.9 times that of whites.


## Congenital syphilis

- CASES: New Hampshire reported 0 cases in 2010.
- RATES: New Hampshire had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, New Hampshire, 2001-2010



## Total P&S Syphilis Rates by County, New Hampshire, 2010



Rate (per 100,000 population)



# New Jersey, 2010

## Primary and secondary syphilis

- CASES: New Jersey reported 244 cases in 2010.
- RATES: New Jersey ranked 27 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.8 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In New Jersey, the rate among males was 5.3 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.4 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In New Jersey, the race/ethnicity adjusted rates per 100,000 population were 1.4 among whites, 8.4 among blacks, 4.5 among Hispanics, 0.4 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 6 times that of whites.
- The rate among Hispanics was 3.2 times that of whites.


## Congenital syphilis

- CASES: New Jersey reported 3 cases in 2010.
- RATES: New Jersey ranked 22 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 2.6 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, New Jersey, 2001-2010



## Total P&S Syphilis Rates by County, New Jersey, 2010



Rate (per 100,000 population)



# New Mexico, 2010

## Primary and secondary syphilis

- CASES: New Mexico reported 53 cases in 2010.
- RATES: New Mexico ranked 32 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.6 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In New Mexico, the rate among males was 5.0 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.3 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In New Mexico, the race/ethnicity adjusted rates per 100,000 population were 1.0 among whites, 4.0 among blacks, 2.5 among Hispanics, 0.0 among Asian/Pacific Islanders, and 5.6 among American Indians/Alaska Natives.
- The rate among blacks was 4 times that of whites.
- The rate among Hispanics was 2.5 times that of whites.
- The rate among American Indians/Alaska Natives was 5.6 times that of whites.


## Congenital syphilis

- CASES: New Mexico reported 0 cases in 2010.
- RATES: New Mexico had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, New Mexico, 2001-2010



## Total P&S Syphilis Rates by County, New Mexico, 2010



Rate (per 100,000 population)



# New York, 2010

## Primary and secondary syphilis

- CASES: New York reported 1,098 cases in 2010.
- RATES: New York ranked 10 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 5.6 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In New York, the rate among males was 11.1 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.5 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In New York, the race/ethnicity adjusted rates per 100,000 population were 2.5 among whites, 14.7 among blacks, 8.9 among Hispanics, 1.1 among Asian/Pacific Islanders, and 2.7 among American Indians/Alaska Natives.
- The rate among blacks was 5.9 times that of whites.
- The rate among Hispanics was 3.6 times that of whites.
- The rate among American Indians/Alaska Natives was 1.1 times that of whites.


## Congenital syphilis

- CASES: New York reported 16 cases in 2010.
- RATES: New York ranked 18 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 6.3 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, New York, 2001-2010



## Total P&S Syphilis Rates by County, New York, 2010



# North Carolina, 2010

## Primary and secondary syphilis

- CASES: North Carolina reported 396 cases in 2010.
- RATES: North Carolina ranked 18 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 4.2 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In North Carolina, the rate among males was 7.4 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 1.1 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In North Carolina, the race/ethnicity adjusted rates per 100,000 population were 0.9 among whites, 15.8 among blacks, 1.7 among Hispanics, 0.5 among Asian/Pacific Islanders, and 0.9 among American Indians/Alaska Natives.
- The rate among blacks was 17.6 times that of whites.
- The rate among Hispanics was 1.9 times that of whites.


## Congenital syphilis

- CASES: North Carolina reported 10 cases in 2010.
- RATES: North Carolina ranked 15 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 7.6 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, North Carolina, 2001-2010



## Total P&S Syphilis Rates by County, North Carolina, 2010



Rate (per 100,000 population)



# North Dakota, 2010

## Primary and secondary syphilis

- CASES: North Dakota reported 3 cases in 2010.
- RATES: North Dakota ranked 48 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 0.5 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In North Dakota, the rate among males was 0.6 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.3 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In North Dakota, the race/ethnicity adjusted rates per 100,000 population were 0.0 among whites, 23.5 among blacks, 0.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.


## Congenital syphilis

- CASES: North Dakota reported 0 cases in 2010.
- RATES: North Dakota had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, North Dakota, 2001-2010



## Total P&S Syphilis Rates by County, North Dakota, 2010



Rate (per 100,000 population)



# Ohio, 2010

## Primary and secondary syphilis

- CASES: Ohio reported 528 cases in 2010.
- RATES: Ohio ranked 15 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 4.6 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Ohio, the rate among males was 7.0 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 2.2 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Ohio, the race/ethnicity adjusted rates per 100,000 population were 1.7 among whites, 24.5 among blacks, 2.5 among Hispanics, 0.5 among Asian/Pacific Islanders, and 3.3 among American Indians/Alaska Natives.
- The rate among blacks was 14.4 times that of whites.
- The rate among Hispanics was 1.5 times that of whites.
- The rate among American Indians/Alaska Natives was 1.9 times that of whites.


## Congenital syphilis

- CASES: Ohio reported 10 cases in 2010.
- RATES: Ohio ranked 17 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 6.6 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Ohio, 2001-2010



## Total P&S Syphilis Rates by County, Ohio, 2010



# Oklahoma, 2010

## Primary and secondary syphilis

- **CASES:** Oklahoma reported 92 cases in 2010.
- **RATES:** Oklahoma ranked 34 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.5 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Oklahoma, the rate among males was 4.2 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.9 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Oklahoma, the race/ethnicity adjusted rates per 100,000 population were 1.6 among whites, 11.5 among blacks, 3.0 among Hispanics, 1.4 among Asian/Pacific Islanders, and 1.3 among American Indians/Alaska Natives.
- The rate among blacks was 7.2 times that of whites.
- The rate among Hispanics was 1.9 times that of whites.


## Congenital syphilis

- **CASES:** Oklahoma reported 0 cases in 2010.
- **RATES:** Oklahoma had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Oklahoma, 2001-2010



## Total P&S Syphilis Rates by County, Oklahoma, 2010



Rate (per 100,000 population)



# Oregon, 2010

## Primary and secondary syphilis

- CASES: Oregon reported 71 cases in 2010.
- RATES: Oregon ranked 38 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 1.9 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Oregon, the rate among males was 3.7 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.1 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Oregon, the race/ethnicity adjusted rates per 100,000 population were 1.8 among whites, 3.6 among blacks, 2.1 among Hispanics, 0.6 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 2 times that of whites.
- The rate among Hispanics was 1.2 times that of whites.


## Congenital syphilis

- CASES: Oregon reported 0 cases in 2010.
- RATES: Oregon had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Oregon, 2001-2010



## Total P&S Syphilis Rates by County, Oregon, 2010



Rate (per 100,000 population)



# Pennsylvania, 2010

## Primary and secondary syphilis

- CASES: Pennsylvania reported 369 cases in 2010.
- RATES: Pennsylvania ranked 25 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.9 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Pennsylvania, the rate among males was 5.4 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.6 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Pennsylvania, the race/ethnicity adjusted rates per 100,000 population were 1.0 among whites, 16.5 among blacks, 3.9 among Hispanics, 0.9 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 16.5 times that of whites.
- The rate among Hispanics was 3.9 times that of whites.


## Congenital syphilis

- CASES: Pennsylvania reported 3 cases in 2010.
- RATES: Pennsylvania ranked 24 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 2.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Pennsylvania, 2001-2010



## Total P&S Syphilis Rates by County, Pennsylvania, 2010



# Rhode Island, 2010

## Primary and secondary syphilis

- **CASES:** Rhode Island reported 41 cases in 2010.
- **RATES:** Rhode Island ranked 20 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 3.9 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Rhode Island, the rate among males was 7.6 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.4 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Rhode Island, the race/ethnicity adjusted rates per 100,000 population were 3.1 among whites, 8.5 among blacks, 7.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 2.7 times that of whites.
- The rate among Hispanics was 2.3 times that of whites.

## Congenital syphilis

- **CASES:** Rhode Island reported 0 cases in 2010.
- **RATES:** Rhode Island had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Rhode Island, 2001-2010



## Total P&S Syphilis Rates by County, Rhode Island, 2010



# South Carolina, 2010

## Primary and secondary syphilis

- CASES: South Carolina reported 155 cases in 2010.
- RATES: South Carolina ranked 23 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 3.4 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In South Carolina, the rate among males was 6.6 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.4 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In South Carolina, the race/ethnicity adjusted rates per 100,000 population were 0.8 among whites, 9.7 among blacks, 1.5 among Hispanics, 1.5 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 12.1 times that of whites.
- The rate among Hispanics was 1.9 times that of whites.
- The rate among Asian/Pacific Islanders was 1.9 times that of whites.


## Congenital syphilis

- CASES: South Carolina reported 0 cases in 2010.
- RATES: South Carolina had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, South Carolina, 2001-2010



## Total P&S Syphilis Rates by County, South Carolina, 2010



Rate (per 100,000 population)



# South Dakota, 2010

## Primary and secondary syphilis

- **CASES:** South Dakota reported 4 cases in 2010.
- **RATES:** South Dakota ranked 47 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 0.5 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In South Dakota, the rate among males was 1.0 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.0 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In South Dakota, the race/ethnicity adjusted rates per 100,000 population were 0.4 among whites, 8.9 among blacks, 0.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 22.3 times that of whites.


## Congenital syphilis

- **CASES:** South Dakota reported 0 cases in 2010.
- **RATES:** South Dakota had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, South Dakota, 2001-2010



## Total P&S Syphilis Rates by County, South Dakota, 2010



Rate (per 100,000 population)



# Tennessee, 2010

## Primary and secondary syphilis

- **CASES:** Tennessee reported 277 cases in 2010.
- **RATES:** Tennessee ranked 16 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 4.4 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Tennessee, the rate among males was 7.4 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 1.5 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Tennessee, the race/ethnicity adjusted rates per 100,000 population were 1.5 among whites, 17.7 among blacks, 3.1 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 11.8 times that of whites.
- The rate among Hispanics was 2.1 times that of whites.


## Congenital syphilis

- **CASES:** Tennessee reported 11 cases in 2010.
- **RATES:** Tennessee ranked 10 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 12.7 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Tennessee, 2001-2010



## Total P&S Syphilis Rates by County, Tennessee, 2010



Rate (per 100,000 population)



# Texas, 2010

## Primary and secondary syphilis

- CASES: Texas reported 1,230 cases in 2010.
- RATES: Texas ranked 13 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 5 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Texas, the rate among males was 7.2 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 2.7 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Texas, the race/ethnicity adjusted rates per 100,000 population were 1.8 among whites, 24.3 among blacks, 3.3 among Hispanics, 1.2 among Asian/Pacific Islanders, and 0.9 among American Indians/Alaska Natives.
- The rate among blacks was 13.5 times that of whites.
- The rate among Hispanics was 1.8 times that of whites.


## Congenital syphilis

- CASES: Texas reported 103 cases in 2010.
- RATES: Texas ranked 4 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 25.3 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Texas, 2001-2010



## Total P&S Syphilis Rates by County, Texas, 2010



Rate (per 100,000 population)



# Utah, 2010

## Primary and secondary syphilis

- CASES: Utah reported 65 cases in 2010.
- RATES: Utah ranked 37 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.3 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Utah, the rate among males was 4.5 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.1 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Utah, the race/ethnicity adjusted rates per 100,000 population were 2.4 among whites, 5.5 among blacks, 2.3 among Hispanics, 1.2 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 2.3 times that of whites.


## Congenital syphilis

- CASES: Utah reported 1 case in 2010.
- RATES: Utah ranked 25 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 1.8 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Utah, 2001-2010



## Total P&S Syphilis Rates by County, Utah, 2010



Rate (per 100,000 population)



# Vermont, 2010

## Primary and secondary syphilis

- CASES: Vermont reported 4 cases in 2010.
- RATES: Vermont ranked 44 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 0.6 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Vermont, the rate among males was 0.7 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.6 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Vermont, the race/ethnicity adjusted rates per 100,000 population were 0.2 among whites, 29.2 among blacks, 0.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 36.9 among American Indians/Alaska Natives.
- The rate among blacks was 146 times that of whites.
- The rate among American Indians/Alaska Natives was 184.5 times that of whites.


## Congenital syphilis

- CASES: Vermont reported 0 cases in 2010.
- RATES: Vermont had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Vermont, 2001-2010



## Total P&S Syphilis Rates by County, Vermont, 2010



Rate (per 100,000 population)



# Virginia, 2010

## Primary and secondary syphilis

- CASES: Virginia reported 279 cases in 2010.
- RATES: Virginia ranked 21 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 3.5 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Virginia, the rate among males was 6.7 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.5 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Virginia, the race/ethnicity adjusted rates per 100,000 population were 1.7 among whites, 11.2 among blacks, 2.3 among Hispanics, 0.2 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 6.6 times that of whites.
- The rate among Hispanics was 1.4 times that of whites.


## Congenital syphilis

- CASES: Virginia reported 1 case in 2010.
- RATES: Virginia ranked 30 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 0.9 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Virginia, 2001-2010



## Total P&S Syphilis Rates by County, Virginia, 2010



Rate (per 100,000 population)



# Washington, 2010

## Primary and secondary syphilis

- CASES: Washington reported 266 cases in 2010.
- RATES: Washington ranked 19 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 4 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Washington, the rate among males was 7.8 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.1 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Washington, the race/ethnicity adjusted rates per 100,000 population were 2.9 among whites, 6.9 among blacks, 4.8 among Hispanics, 1.7 among Asian/Pacific Islanders, and 0.9 among American Indians/Alaska Natives.
- The rate among blacks was 2.4 times that of whites.
- The rate among Hispanics was 1.7 times that of whites.


## Congenital syphilis

- CASES: Washington reported 1 case in 2010.
- RATES: Washington ranked 29 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 1.1 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Washington, 2001-2010



## Total P&S Syphilis Rates by County, Washington, 2010



# West Virginia, 2010

## Primary and secondary syphilis

- **CASES:** West Virginia reported 6 cases in 2010.
- **RATES:** West Virginia ranked 51 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 0.3 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In West Virginia, the rate among males was 0.7 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.0 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In West Virginia, the race/ethnicity adjusted rates per 100,000 population were 0.2 among whites, 0.0 among blacks, 0.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.


## Congenital syphilis

- **CASES:** West Virginia reported 0 cases in 2010.
- **RATES:** West Virginia had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, West Virginia, 2001-2010



## Total P&S Syphilis Rates by County, West Virginia, 2010



# Wisconsin, 2010

## Primary and secondary syphilis

- CASES: Wisconsin reported 49 cases in 2010.
- RATES: Wisconsin ranked 41 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 0.9 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Wisconsin, the rate among males was 1.5 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.2 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Wisconsin, the race/ethnicity adjusted rates per 100,000 population were 0.3 among whites, 5.9 among blacks, 1.7 among Hispanics, 1.6 among Asian/Pacific Islanders, and 1.9 among American Indians/Alaska Natives.
- The rate among blacks was 19.7 times that of whites.
- The rate among Hispanics was 5.7 times that of whites.
- The rate among Asian/Pacific Islanders was 5.3 times that of whites.
- The rate among American Indians/Alaska Natives was 6.3 times that of whites.


## Congenital syphilis

- CASES: Wisconsin reported 1 case in 2010.
- RATES: Wisconsin ranked 27 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 1.4 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Wisconsin, 2001-2010



## Total P&S Syphilis Rates by County, Wisconsin, 2010



# Wyoming, 2010

## Primary and secondary syphilis

- CASES: Wyoming reported 0 cases in 2010.
- RATES: Wyoming reported 0 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Wyoming, the rate among males was 0.0 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.0 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Wyoming, the race/ethnicity adjusted rates per 100,000 population were 0.0 among whites, 0.0 among blacks, 0.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.


## Congenital syphilis

- CASES: Wyoming reported 0 cases in 2010.
- RATES: Wyoming had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Wyoming, 2001-2010



## Total P&S Syphilis Rates by County, Wyoming, 2010



Rate (per 100,000 population)



# Guam, 2010

## Primary and secondary syphilis

- CASES: Guam reported 1 case in 2010.
- RATES: Guam ranked 46 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 0.6 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.


## Rates of primary and secondary syphilis, by sex

- In Guam, the rate among males was 0.0 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 1.1 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Congenital syphilis

- CASES: Guam reported 0 cases in 2010.
- RATES: Guam had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Guam, 2001-2010



## Total P&S Syphilis Rate, Guam, 2010



# Puerto Rico, 2010

## Primary and secondary syphilis

- CASES: Puerto Rico reported 228 cases in 2010.
- RATES: Puerto Rico ranked 9 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 5.7 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.


## Rates of primary and secondary syphilis, by sex

- In Puerto Rico, the rate among males was 11.0 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.9 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Congenital syphilis

- CASES: Puerto Rico reported 2 cases in 2010.
- RATES: Puerto Rico ranked 20 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 4.3 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Puerto Rico, 2001-2010



## Total P&S Syphilis Rates by Municipality, Puerto Rico, 2010



Rate (per 100,000 population)



# Virgin Islands, 2010

## Primary and secondary syphilis

- CASES: Virgin Islands reported 0 cases in 2010.
- RATES: Virgin Islands reported 0 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.


## Rates of primary and secondary syphilis, by sex

- In Virgin Islands, the rate among males was 0.0 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.0 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.

## Congenital syphilis

- CASES: Virgin Islands reported 0 cases in 2010.
- RATES: Virgin Islands had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Virgin Islands, 2001-2010



Total P&S Syphilis Rate, Virgin Islands, 2010



Rate (per 100,000 population)

