

Connecticut, 2011

Primary and secondary syphilis

- **CASES:** Connecticut reported 65 cases in 2011.
- **RATES:** Connecticut ranked 37 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 1.8 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Connecticut, the rate among males was 3.4 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.3 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Connecticut, the race/ethnicity adjusted rates per 100,000 population were 1.0 among whites, 5.6 among blacks, 3.8 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 5.6 times that of whites.
- The rate among Hispanics was 3.8 times that of whites.

Congenital syphilis

- **CASES:** Connecticut reported 0 cases in 2011.
- **RATES:** Connecticut had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Connecticut, 2002 – 2011



Total P&S Syphilis Rates by County, Connecticut, 2011

