

District of Columbia, 2011

Primary and secondary syphilis

- **CASES:** District of Columbia reported 165 cases in 2011.
- **RATES:** District of Columbia ranked 1 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 27.4 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In District of Columbia, the rate among males was 55.6 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 2.2 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In District of Columbia, the race/ethnicity adjusted rates per 100,000 population were 19.6 among whites, 33.1 among blacks, 23.7 among Hispanics, 0.0 among Asian/Pacific Islanders, and 57.9 among American Indians/Alaska Natives.
- The rate among blacks was 1.7 times that of whites.
- The rate among Hispanics was 1.2 times that of whites.
- The rate among American Indians/Alaska Natives was 3 times that of whites.


Congenital syphilis

- **CASES:** District of Columbia reported 1 case in 2011.
- **RATES:** District of Columbia ranked 9 among among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 11.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, District of Columbia, 2002 – 2011



Total P&S Syphilis Rate, District of Columbia, 2011



Rate (per 100,000 population)

