

Indiana, 2011

Primary and secondary syphilis

- CASES: Indiana reported 173 cases in 2011.
- RATES: Indiana ranked 29 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.7 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Indiana, the rate among males was 5.0 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.4 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Indiana, the race/ethnicity adjusted rates per 100,000 population were 1.5 among whites, 13.0 among blacks, 2.3 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 8.7 times that of whites.
- The rate among Hispanics was 1.5 times that of whites.


Congenital syphilis

- CASES: Indiana reported 0 cases in 2011.
- RATES: Indiana had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Indiana, 2002 – 2011



Total P&S Syphilis Rates by County, Indiana, 2011



Rate (per 100,000 population)

