

Ohio, 2011

Primary and secondary syphilis

- CASES: Ohio reported 440 cases in 2011.
- RATES: Ohio ranked 22 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 3.8 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Ohio, the rate among males was 5.9 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 1.8 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Ohio, the race/ethnicity adjusted rates per 100,000 population were 1.4 among whites, 19.7 among blacks, 1.7 among Hispanics, 0.0 among Asian/Pacific Islanders, and 3.5 among American Indians/Alaska Natives.
- The rate among blacks was 14.1 times that of whites.
- The rate among Hispanics was 1.2 times that of whites.
- The rate among American Indians/Alaska Natives was 2.5 times that of whites.

Congenital syphilis

- CASES: Ohio reported 13 cases in 2011.
- RATES: Ohio ranked 12 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 8.7 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Ohio, 2002 – 2011



Total P&S Syphilis Rates by County, Ohio, 2011

