

Alabama, 2011

Primary and secondary syphilis

- CASES: Alabama reported 228 cases in 2011.
- RATES: Alabama ranked 15 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 4.8 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Alabama, the rate among males was 7.5 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 2.2 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Alabama, the race/ethnicity adjusted rates per 100,000 population were 1.9 among whites, 13.0 among blacks, 1.1 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 6.8 times that of whites.


Congenital syphilis

- CASES: Alabama reported 10 cases in 2011.
- RATES: Alabama ranked 5 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 15.5 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Alabama, 2002 – 2011



Total P&S Syphilis Rates by County, Alabama, 2011



Rate (per 100,000 population)



Alaska, 2011

Primary and secondary syphilis

- CASES: Alaska reported 5 cases in 2011.
- RATES: Alaska ranked 46 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 0.7 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Alaska, the rate among males was 1.1 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.3 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Alaska, the race/ethnicity adjusted rates per 100,000 population were 0.4 among whites, 3.5 among blacks, 2.5 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.9 among American Indians/Alaska Natives.
- The rate among blacks was 8.8 times that of whites.
- The rate among Hispanics was 6.3 times that of whites.
- The rate among American Indians/Alaska Natives was 2.3 times that of whites.


Congenital syphilis

- CASES: Alaska reported 0 cases in 2011.
- RATES: Alaska had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Alaska, 2002 – 2011



Total P&S Syphilis Rates by County, Alaska, 2011



Arizona, 2011

Primary and secondary syphilis

- CASES: Arizona reported 274 cases in 2011.
- RATES: Arizona ranked 20 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 4.3 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Arizona, the rate among males was 8.1 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.5 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Arizona, the race/ethnicity adjusted rates per 100,000 population were 2.9 among whites, 14.5 among blacks, 5.6 among Hispanics, 2.0 among Asian/Pacific Islanders, and 4.8 among American Indians/Alaska Natives.
- The rate among blacks was 5 times that of whites.
- The rate among Hispanics was 1.9 times that of whites.
- The rate among American Indians/Alaska Natives was 1.7 times that of whites.


Congenital syphilis

- CASES: Arizona reported 14 cases in 2011.
- RATES: Arizona ranked 6 among among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 14.1 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Arizona, 2002 – 2011



Total P&S Syphilis Rates by County, Arizona, 2011



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Arkansas, 2011

Primary and secondary syphilis

- CASES: Arkansas reported 182 cases in 2011.
- RATES: Arkansas ranked 10 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 6.2 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Arkansas, the rate among males was 7.4 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 5.1 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Arkansas, the race/ethnicity adjusted rates per 100,000 population were 1.7 among whites, 30.5 among blacks, 1.6 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 17.9 times that of whites.


Congenital syphilis

- CASES: Arkansas reported 15 cases in 2011.
- RATES: Arkansas ranked 1 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 36.9 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Arkansas, 2002 – 2011



Total P&S Syphilis Rates by County, Arkansas, 2011



Rate (per 100,000 population)



California, 2011

Primary and secondary syphilis

- CASES: California reported 2,443 cases in 2011.
- RATES: California ranked 8 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 6.6 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In California, the rate among males was 12.6 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.5 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In California, the race/ethnicity adjusted rates per 100,000 population were 6.5 among whites, 16.2 among blacks, 5.7 among Hispanics, 2.2 among Asian/Pacific Islanders, and 5.8 among American Indians/Alaska Natives.
- The rate among blacks was 2.5 times that of whites.


Congenital syphilis

- CASES: California reported 40 cases in 2011.
- RATES: California ranked 14 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 7.2 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, California, 2002 – 2011



Total P&S Syphilis Rates by County, California, 2011



Rate (per 100,000 population)

≤0.4 0.41 - 2.2 >2.2

Colorado, 2011

Primary and secondary syphilis

- CASES: Colorado reported 133 cases in 2011.
- RATES: Colorado ranked 31 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.6 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Colorado, the rate among males was 5.1 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.2 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Colorado, the race/ethnicity adjusted rates per 100,000 population were 2.2 among whites, 5.1 among blacks, 3.7 among Hispanics, 1.9 among Asian/Pacific Islanders, and 8.0 among American Indians/Alaska Natives.
- The rate among blacks was 2.3 times that of whites.
- The rate among Hispanics was 1.7 times that of whites.
- The rate among American Indians/Alaska Natives was 3.6 times that of whites.


Congenital syphilis

- CASES: Colorado reported 0 cases in 2011.
- RATES: Colorado had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Colorado, 2002 – 2011



Total P&S Syphilis Rates by County, Colorado, 2011



Connecticut, 2011

Primary and secondary syphilis

- CASES: Connecticut reported 65 cases in 2011.
- RATES: Connecticut ranked 37 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 1.8 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Connecticut, the rate among males was 3.4 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.3 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Connecticut, the race/ethnicity adjusted rates per 100,000 population were 1.0 among whites, 5.6 among blacks, 3.8 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 5.6 times that of whites.
- The rate among Hispanics was 3.8 times that of whites.


Congenital syphilis

- CASES: Connecticut reported 0 cases in 2011.
- RATES: Connecticut had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Connecticut, 2002 – 2011



Total P&S Syphilis Rates by County, Connecticut, 2011



Delaware, 2011

Primary and secondary syphilis

- CASES: Delaware reported 27 cases in 2011.
- RATES: Delaware ranked 24 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 3.0 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Delaware, the rate among males was 6.0 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.2 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Delaware, the race/ethnicity adjusted rates per 100,000 population were 1.8 among whites, 8.2 among blacks, 0.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 4.6 times that of whites.


Congenital syphilis

- CASES: Delaware reported 0 cases in 2011.
- RATES: Delaware had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Delaware, 2002 – 2011



Total P&S Syphilis Rates by County, Delaware, 2011



Rate (per 100,000 population)

<=0.4

0.41 - 2.2

>2.2

District of Columbia, 2011

Primary and secondary syphilis

- CASES: District of Columbia reported 165 cases in 2011.
- RATES: District of Columbia ranked 1 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 27.4 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In District of Columbia, the rate among males was 55.6 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 2.2 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In District of Columbia, the race/ethnicity adjusted rates per 100,000 population were 19.6 among whites, 33.1 among blacks, 23.7 among Hispanics, 0.0 among Asian/Pacific Islanders, and 57.9 among American Indians/Alaska Natives.
- The rate among blacks was 1.7 times that of whites.
- The rate among Hispanics was 1.2 times that of whites.
- The rate among American Indians/Alaska Natives was 3 times that of whites.


Congenital syphilis

- CASES: District of Columbia reported 1 case in 2011.
- RATES: District of Columbia ranked 9 among among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 11.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, District of Columbia, 2002 – 2011



Total P&S Syphilis Rate, District of Columbia, 2011



Rate (per 100,000 population)

<=0.4

0.41 - 2.2

>2.2

Florida, 2011

Primary and secondary syphilis

- CASES: Florida reported 1,257 cases in 2011.
- RATES: Florida ranked 7 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 6.7 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Florida, the rate among males was 12.2 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 1.4 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Florida, the race/ethnicity adjusted rates per 100,000 population were 4.2 among whites, 15.6 among blacks, 6.5 among Hispanics, 1.6 among Asian/Pacific Islanders, and 1.8 among American Indians/Alaska Natives.
- The rate among blacks was 3.7 times that of whites.
- The rate among Hispanics was 1.5 times that of whites.


Congenital syphilis

- CASES: Florida reported 32 cases in 2011.
- RATES: Florida ranked 7 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 13.8 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Florida, 2002 – 2011



Total P&S Syphilis Rates by County, Florida, 2011



Rate (per 100,000 population)

<=0.4

0.41 - 2.2

>2.2

Georgia, 2011

Primary and secondary syphilis

- CASES: Georgia reported 678 cases in 2011.
- RATES: Georgia ranked 4 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 7.0 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Georgia, the rate among males was 13.1 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 1.2 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Georgia, the race/ethnicity adjusted rates per 100,000 population were 1.9 among whites, 17.7 among blacks, 2.0 among Hispanics, 0.9 among Asian/Pacific Islanders, and 12.2 among American Indians/Alaska Natives.
- The rate among blacks was 9.3 times that of whites.
- The rate among Hispanics was 1.1 times that of whites.
- The rate among American Indians/Alaska Natives was 6.4 times that of whites.


Congenital syphilis

- CASES: Georgia reported 10 cases in 2011.
- RATES: Georgia ranked 15 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 6.8 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Georgia, 2002 – 2011



Total P&S Syphilis Rates by County, Georgia, 2011



Rate (per 100,000 population)



Hawaii, 2011

Primary and secondary syphilis

- CASES: Hawaii reported 14 cases in 2011.
- RATES: Hawaii ranked 41 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 1.0 case per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Hawaii, the rate among males was 2.1 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.0 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Hawaii, the race/ethnicity adjusted rates per 100,000 population were 2.5 among whites, 6.9 among blacks, 0.0 among Hispanics, 0.1 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 2.8 times that of whites.


Congenital syphilis

- CASES: Hawaii reported 0 cases in 2011.
- RATES: Hawaii had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Hawaii, 2002 – 2011



Total P&S Syphilis Rates by County, Hawaii, 2011



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Idaho, 2011

Primary and secondary syphilis

- CASES: Idaho reported 13 cases in 2011.
- RATES: Idaho ranked 44 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 0.8 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Idaho, the rate among males was 1.5 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.1 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Idaho, the race/ethnicity adjusted rates per 100,000 population were 0.5 among whites, 0.0 among blacks, 1.1 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among Hispanics was 2.2 times that of whites.


Congenital syphilis

- CASES: Idaho reported 0 cases in 2011.
- RATES: Idaho had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Idaho, 2002 – 2011



Total P&S Syphilis Rates by County, Idaho, 2011



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Illinois, 2011

Primary and secondary syphilis

- CASES: Illinois reported 881 cases in 2011.
- RATES: Illinois ranked 5 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 6.9 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Illinois, the rate among males was 12.7 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 1.2 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Illinois, the race/ethnicity adjusted rates per 100,000 population were 2.9 among whites, 25.4 among blacks, 5.7 among Hispanics, 1.5 among Asian/Pacific Islanders, and 4.0 among American Indians/Alaska Natives.
- The rate among blacks was 8.8 times that of whites.
- The rate among Hispanics was 2 times that of whites.
- The rate among American Indians/Alaska Natives was 1.4 times that of whites.


Congenital syphilis

- CASES: Illinois reported 18 cases in 2011.
- RATES: Illinois ranked 10 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 10.2 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Illinois, 2002 – 2011



Total P&S Syphilis Rates by County, Illinois, 2011



Rate (per 100,000 population)



Indiana, 2011

Primary and secondary syphilis

- CASES: Indiana reported 173 cases in 2011.
- RATES: Indiana ranked 29 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.7 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Indiana, the rate among males was 5.0 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.4 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Indiana, the race/ethnicity adjusted rates per 100,000 population were 1.5 among whites, 13.0 among blacks, 2.3 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 8.7 times that of whites.
- The rate among Hispanics was 1.5 times that of whites.


Congenital syphilis

- CASES: Indiana reported 0 cases in 2011.
- RATES: Indiana had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Indiana, 2002 – 2011



Total P&S Syphilis Rates by County, Indiana, 2011



Rate (per 100,000 population)

≤ 0.4
 0.41 - 2.2
 > 2.2

Iowa, 2011

Primary and secondary syphilis

- CASES: Iowa reported 20 cases in 2011.
- RATES: Iowa ranked 47 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 0.7 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Iowa, the rate among males was 1.0 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.3 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Iowa, the race/ethnicity adjusted rates per 100,000 population were 0.5 among whites, 3.9 among blacks, 0.7 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 7.8 times that of whites.
- The rate among Hispanics was 1.4 times that of whites.


Congenital syphilis

- CASES: Iowa reported 0 cases in 2011.
- RATES: Iowa had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Iowa, 2002 – 2011



Total P&S Syphilis Rates by County, Iowa, 2011



Rate (per 100,000 population)



Kansas, 2011

Primary and secondary syphilis

- CASES: Kansas reported 24 cases in 2011.
- RATES: Kansas ranked 43 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 0.8 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Kansas, the rate among males was 1.7 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.0 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Kansas, the race/ethnicity adjusted rates per 100,000 population were 0.5 among whites, 4.9 among blacks, 1.0 among Hispanics, 1.3 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 9.8 times that of whites.
- The rate among Hispanics was 2 times that of whites.
- The rate among Asian/Pacific Islanders was 2.6 times that of whites.


Congenital syphilis

- CASES: Kansas reported 0 cases in 2011.
- RATES: Kansas had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Kansas, 2002 – 2011



Total P&S Syphilis Rates by County, Kansas, 2011



Rate (per 100,000 population)

≤ 0.4
 0.41 - 2.2
 > 2.2

Kentucky, 2011

Primary and secondary syphilis

- CASES: Kentucky reported 129 cases in 2011.
- RATES: Kentucky ranked 25 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 3.0 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Kentucky, the rate among males was 5.2 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.9 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Kentucky, the race/ethnicity adjusted rates per 100,000 population were 1.9 among whites, 14.1 among blacks, 3.8 among Hispanics, 1.8 among Asian/Pacific Islanders, and 10.8 among American Indians/Alaska Natives.
- The rate among blacks was 7.4 times that of whites.
- The rate among Hispanics was 2 times that of whites.
- The rate among American Indians/Alaska Natives was 5.7 times that of whites.


Congenital syphilis

- CASES: Kentucky reported 2 cases in 2011.
- RATES: Kentucky ranked 22 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 3.4 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Kentucky, 2002 – 2011



Total P&S Syphilis Rates by County, Kentucky, 2011



Rate (per 100,000 population)

≤0.4 0.41 - 2.2 >2.2

Louisiana, 2011

Primary and secondary syphilis

- CASES: Louisiana reported 447 cases in 2011.
- RATES: Louisiana ranked 2 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 9.9 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Louisiana, the rate among males was 12.1 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 7.7 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Louisiana, the race/ethnicity adjusted rates per 100,000 population were 2.5 among whites, 25.2 among blacks, 2.1 among Hispanics, 1.3 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 10.1 times that of whites.


Congenital syphilis

- CASES: Louisiana reported 18 cases in 2011.
- RATES: Louisiana ranked 3 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 27.6 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Louisiana, 2002 – 2011



Total P&S Syphilis Rates by County, Louisiana, 2011



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Maine, 2011

Primary and secondary syphilis

- CASES: Maine reported 12 cases in 2011.
- RATES: Maine ranked 42 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 0.9 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Maine, the rate among males was 1.8 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.0 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Maine, the race/ethnicity adjusted rates per 100,000 population were 0.7 among whites, 0.0 among blacks, 0.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.


Congenital syphilis

- CASES: Maine reported 0 cases in 2011.
- RATES: Maine had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Maine, 2002 – 2011



Total P&S Syphilis Rates by County, Maine, 2011



Rate (per 100,000 population)



Maryland, 2011

Primary and secondary syphilis

- CASES: Maryland reported 452 cases in 2011.
- RATES: Maryland ranked 3 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 7.8 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Maryland, the rate among males was 14.4 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 1.6 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Maryland, the race/ethnicity adjusted rates per 100,000 population were 2.0 among whites, 20.9 among blacks, 2.8 among Hispanics, 1.2 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 10.5 times that of whites.
- The rate among Hispanics was 1.4 times that of whites.


Congenital syphilis

- CASES: Maryland reported 24 cases in 2011.
- RATES: Maryland ranked 2 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 31.1 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Maryland, 2002 – 2011



Total P&S Syphilis Rates by County, Maryland, 2011



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Massachusetts, 2011

Primary and secondary syphilis

- CASES: Massachusetts reported 266 cases in 2011.
- RATES: Massachusetts ranked 21 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 4.1 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Massachusetts, the rate among males was 7.7 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.7 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Massachusetts, the race/ethnicity adjusted rates per 100,000 population were 2.5 among whites, 9.7 among blacks, 10.2 among Hispanics, 3.0 among Asian/Pacific Islanders, and 6.8 among American Indians/Alaska Natives.
- The rate among blacks was 3.9 times that of whites.
- The rate among Hispanics was 4.1 times that of whites.
- The rate among Asian/Pacific Islanders was 1.2 times that of whites.
- The rate among American Indians/Alaska Natives was 2.7 times that of whites.


Congenital syphilis

- CASES: Massachusetts reported 0 cases in 2011.
- RATES: Massachusetts had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Massachusetts, 2002 – 2011



Total P&S Syphilis Rates by County, Massachusetts, 2011



Rate (per 100,000 population)

<=0.4

0.41 - 2.2

>2.2

Michigan, 2011

Primary and secondary syphilis

- CASES: Michigan reported 286 cases in 2011.
- RATES: Michigan ranked 27 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.9 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Michigan, the rate among males was 5.4 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.5 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Michigan, the race/ethnicity adjusted rates per 100,000 population were 1.2 among whites, 12.5 among blacks, 0.9 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 10.4 times that of whites.


Congenital syphilis

- CASES: Michigan reported 6 cases in 2011.
- RATES: Michigan ranked 17 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 5.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Michigan, 2002 – 2011



Total P&S Syphilis Rates by County, Michigan, 2011



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Minnesota, 2011

Primary and secondary syphilis

- CASES: Minnesota reported 139 cases in 2011.
- RATES: Minnesota ranked 33 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.6 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Minnesota, the rate among males was 5.1 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.2 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Minnesota, the race/ethnicity adjusted rates per 100,000 population were 2.2 among whites, 8.4 among blacks, 4.0 among Hispanics, 1.8 among Asian/Pacific Islanders, and 1.6 among American Indians/Alaska Natives.
- The rate among blacks was 3.8 times that of whites.
- The rate among Hispanics was 1.8 times that of whites.


Congenital syphilis

- CASES: Minnesota reported 0 cases in 2011.
- RATES: Minnesota had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Minnesota, 2002 – 2011



Total P&S Syphilis Rates by County, Minnesota, 2011



Rate (per 100,000 population)



Mississippi, 2011

Primary and secondary syphilis

- CASES: Mississippi reported 191 cases in 2011.
- RATES: Mississippi ranked 9 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 6.4 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Mississippi, the rate among males was 10.1 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 2.9 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Mississippi, the race/ethnicity adjusted rates per 100,000 population were 1.7 among whites, 14.3 among blacks, 0.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 8.4 times that of whites.


Congenital syphilis

- CASES: Mississippi reported 6 cases in 2011.
- RATES: Mississippi ranked 8 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 13.3 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Mississippi, 2002 – 2011



Total P&S Syphilis Rates by County, Mississippi, 2011



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Missouri, 2011

Primary and secondary syphilis

- CASES: Missouri reported 136 cases in 2011.
- RATES: Missouri ranked 35 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.3 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Missouri, the rate among males was 4.4 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.2 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Missouri, the race/ethnicity adjusted rates per 100,000 population were 1.1 among whites, 10.1 among blacks, 1.9 among Hispanics, 1.8 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 9.2 times that of whites.
- The rate among Hispanics was 1.7 times that of whites.
- The rate among Asian/Pacific Islanders was 1.6 times that of whites.


Congenital syphilis

- CASES: Missouri reported 1 case in 2011.
- RATES: Missouri ranked 25 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 1.2 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Missouri, 2002 – 2011



Total P&S Syphilis Rates by County, Missouri, 2011



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Montana, 2011

Primary and secondary syphilis

- CASES: Montana reported 7 cases in 2011.
- RATES: Montana ranked 45 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 0.7 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Montana, the rate among males was 1.2 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.2 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Montana, the race/ethnicity adjusted rates per 100,000 population were 0.6 among whites, 0.0 among blacks, 3.5 among Hispanics, 0.0 among Asian/Pacific Islanders, and 1.5 among American Indians/Alaska Natives.
- The rate among Hispanics was 5.8 times that of whites.
- The rate among American Indians/Alaska Natives was 2.5 times that of whites.


Congenital syphilis

- CASES: Montana reported 0 cases in 2011.
- RATES: Montana had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Montana, 2002 – 2011



Total P&S Syphilis Rates by County, Montana, 2011



Rate (per 100,000 population)

<=0.4

 0.41 - 2.2

 >2.2

Nebraska, 2011

Primary and secondary syphilis

- CASES: Nebraska reported 10 cases in 2011.
- RATES: Nebraska ranked 48 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 0.5 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Nebraska, the rate among males was 1.0 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.1 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Nebraska, the race/ethnicity adjusted rates per 100,000 population were 0.2 among whites, 4.4 among blacks, 0.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 22 times that of whites.


Congenital syphilis

- CASES: Nebraska reported 0 cases in 2011.
- RATES: Nebraska had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Nebraska, 2002 – 2011



Total P&S Syphilis Rates by County, Nebraska, 2011



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Nevada, 2011

Primary and secondary syphilis

- CASES: Nevada reported 136 cases in 2011.
- RATES: Nevada ranked 12 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 5.0 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Nevada, the rate among males was 9.5 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.5 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Nevada, the race/ethnicity adjusted rates per 100,000 population were 3.9 among whites, 13.1 among blacks, 3.5 among Hispanics, 3.6 among Asian/Pacific Islanders, and 7.4 among American Indians/Alaska Natives.
- The rate among blacks was 3.4 times that of whites.
- The rate among American Indians/Alaska Natives was 1.9 times that of whites.


Congenital syphilis

- CASES: Nevada reported 3 cases in 2011.
- RATES: Nevada ranked 13 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 7.6 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Nevada, 2002 – 2011



Total P&S Syphilis Rates by County, Nevada, 2011



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

New Hampshire, 2011

Primary and secondary syphilis

- CASES: New Hampshire reported 18 cases in 2011.
- RATES: New Hampshire ranked 39 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 1.4 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In New Hampshire, the rate among males was 2.6 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.1 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In New Hampshire, the race/ethnicity adjusted rates per 100,000 population were 1.1 among whites, 5.9 among blacks, 8.2 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 5.4 times that of whites.
- The rate among Hispanics was 7.5 times that of whites.


Congenital syphilis

- CASES: New Hampshire reported 0 cases in 2011.
- RATES: New Hampshire had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, New Hampshire, 2002 – 2011



Total P&S Syphilis Rates by County, New Hampshire, 2011



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

New Jersey, 2011

Primary and secondary syphilis

- CASES: New Jersey reported 232 cases in 2011.
- RATES: New Jersey ranked 32 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.6 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In New Jersey, the rate among males was 5.1 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.3 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In New Jersey, the race/ethnicity adjusted rates per 100,000 population were 1.5 among whites, 7.1 among blacks, 3.9 among Hispanics, 0.9 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 4.7 times that of whites.
- The rate among Hispanics was 2.6 times that of whites.


Congenital syphilis

- CASES: New Jersey reported 5 cases in 2011.
- RATES: New Jersey ranked 18 among among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 4.4 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, New Jersey, 2002 – 2011



Total P&S Syphilis Rates by County, New Jersey, 2011



Rate (per 100,000 population)



New Mexico, 2011

Primary and secondary syphilis

- CASES: New Mexico reported 71 cases in 2011.
- RATES: New Mexico ranked 23 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 3.4 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In New Mexico, the rate among males was 6.8 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.2 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In New Mexico, the race/ethnicity adjusted rates per 100,000 population were 1.4 among whites, 7.1 among blacks, 3.4 among Hispanics, 0.0 among Asian/Pacific Islanders, and 6.6 among American Indians/Alaska Natives.
- The rate among blacks was 5.1 times that of whites.
- The rate among Hispanics was 2.4 times that of whites.
- The rate among American Indians/Alaska Natives was 4.7 times that of whites.


Congenital syphilis

- CASES: New Mexico reported 0 cases in 2011.
- RATES: New Mexico had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, New Mexico, 2002 – 2011



Total P&S Syphilis Rates by County, New Mexico, 2011



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

New York, 2011

Primary and secondary syphilis

- CASES: New York reported 1,083 cases in 2011.
- RATES: New York ranked 11 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 5.6 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In New York, the rate among males was 11.1 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.4 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In New York, the race/ethnicity adjusted rates per 100,000 population were 2.4 among whites, 13.7 among blacks, 6.8 among Hispanics, 2.4 among Asian/Pacific Islanders, and 7.2 among American Indians/Alaska Natives.
- The rate among blacks was 5.7 times that of whites.
- The rate among Hispanics was 2.8 times that of whites.
- The rate among American Indians/Alaska Natives was 3 times that of whites.


Congenital syphilis

- CASES: New York reported 13 cases in 2011.
- RATES: New York ranked 16 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 5.2 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, New York, 2002 – 2011



Total P&S Syphilis Rates by County, New York, 2011



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

North Carolina, 2011

Primary and secondary syphilis

- CASES: North Carolina reported 431 cases in 2011.
- RATES: North Carolina ranked 17 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 4.5 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In North Carolina, the rate among males was 8.6 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.6 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In North Carolina, the race/ethnicity adjusted rates per 100,000 population were 1.5 among whites, 15.3 among blacks, 1.4 among Hispanics, 1.8 among Asian/Pacific Islanders, and 0.9 among American Indians/Alaska Natives.
- The rate among blacks was 10.2 times that of whites.
- The rate among Asian/Pacific Islanders was 1.2 times that of whites.


Congenital syphilis

- CASES: North Carolina reported 5 cases in 2011.
- RATES: North Carolina ranked 20 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 3.8 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, North Carolina, 2002 – 2011



Total P&S Syphilis Rates by County, North Carolina, 2011



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

North Dakota, 2011

Primary and secondary syphilis

- CASES: North Dakota reported 1 case in 2011.
- RATES: North Dakota ranked 51 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 0.1 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In North Dakota, the rate among males was 0.3 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.0 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In North Dakota, the race/ethnicity adjusted rates per 100,000 population were 0.0 among whites, 0.0 among blacks, 7.4 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.


Congenital syphilis

- CASES: North Dakota reported 0 cases in 2011.
- RATES: North Dakota had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, North Dakota, 2002 – 2011



Total P&S Syphilis Rates by County, North Dakota, 2011



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Ohio, 2011

Primary and secondary syphilis

- CASES: Ohio reported 440 cases in 2011.
- RATES: Ohio ranked 22 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 3.8 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Ohio, the rate among males was 5.9 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 1.8 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Ohio, the race/ethnicity adjusted rates per 100,000 population were 1.4 among whites, 19.7 among blacks, 1.7 among Hispanics, 0.0 among Asian/Pacific Islanders, and 3.5 among American Indians/Alaska Natives.
- The rate among blacks was 14.1 times that of whites.
- The rate among Hispanics was 1.2 times that of whites.
- The rate among American Indians/Alaska Natives was 2.5 times that of whites.


Congenital syphilis

- CASES: Ohio reported 13 cases in 2011.
- RATES: Ohio ranked 12 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 8.7 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Ohio, 2002 – 2011



Total P&S Syphilis Rates by County, Ohio, 2011



Oklahoma, 2011

Primary and secondary syphilis

- CASES: Oklahoma reported 84 cases in 2011.
- RATES: Oklahoma ranked 36 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.2 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Oklahoma, the rate among males was 3.9 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.6 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Oklahoma, the race/ethnicity adjusted rates per 100,000 population were 1.6 among whites, 8.9 among blacks, 1.2 among Hispanics, 4.0 among Asian/Pacific Islanders, and 1.4 among American Indians/Alaska Natives.
- The rate among blacks was 5.6 times that of whites.
- The rate among Asian/Pacific Islanders was 2.5 times that of whites.


Congenital syphilis

- CASES: Oklahoma reported 2 cases in 2011.
- RATES: Oklahoma ranked 21 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 3.7 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Oklahoma, 2002 – 2011



Total P&S Syphilis Rates by County, Oklahoma, 2011



Rate (per 100,000 population)



Oregon, 2011

Primary and secondary syphilis

- CASES: Oregon reported 97 cases in 2011.
- RATES: Oregon ranked 34 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.5 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Oregon, the rate among males was 5.1 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.0 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Oregon, the race/ethnicity adjusted rates per 100,000 population were 2.5 among whites, 7.3 among blacks, 2.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 1.9 among American Indians/Alaska Natives.
- The rate among blacks was 2.9 times that of whites.


Congenital syphilis

- CASES: Oregon reported 0 cases in 2011.
- RATES: Oregon had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Oregon, 2002 – 2011



Total P&S Syphilis Rates by County, Oregon, 2011



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Pennsylvania, 2011

Primary and secondary syphilis

- CASES: Pennsylvania reported 373 cases in 2011.
- RATES: Pennsylvania ranked 26 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.9 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Pennsylvania, the rate among males was 5.5 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.5 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Pennsylvania, the race/ethnicity adjusted rates per 100,000 population were 1.4 among whites, 12.9 among blacks, 4.7 among Hispanics, 0.3 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 9.2 times that of whites.
- The rate among Hispanics was 3.4 times that of whites.


Congenital syphilis

- CASES: Pennsylvania reported 5 cases in 2011.
- RATES: Pennsylvania ranked 23 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 3.3 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Pennsylvania, 2002 – 2011



Total P&S Syphilis Rates by County, Pennsylvania, 2011



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Rhode Island, 2011

Primary and secondary syphilis

- CASES: Rhode Island reported 46 cases in 2011.
- RATES: Rhode Island ranked 19 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 4.4 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Rhode Island, the rate among males was 8.5 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.6 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Rhode Island, the race/ethnicity adjusted rates per 100,000 population were 3.7 among whites, 14.1 among blacks, 0.8 among Hispanics, 3.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 3.8 times that of whites.


Congenital syphilis

- CASES: Rhode Island reported 0 cases in 2011.
- RATES: Rhode Island had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Rhode Island, 2002 – 2011



Total P&S Syphilis Rates by County, Rhode Island, 2011



Rate (per 100,000 population)

<=0.4

0.41 - 2.2

>2.2

South Carolina, 2011

Primary and secondary syphilis

- CASES: South Carolina reported 221 cases in 2011.
- RATES: South Carolina ranked 14 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 4.8 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In South Carolina, the rate among males was 8.8 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 1.0 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In South Carolina, the race/ethnicity adjusted rates per 100,000 population were 1.5 among whites, 13.1 among blacks, 1.3 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 8.7 times that of whites.


Congenital syphilis

- CASES: South Carolina reported 0 cases in 2011.
- RATES: South Carolina had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, South Carolina, 2002 – 2011



Total P&S Syphilis Rates by County, South Carolina, 2011



Rate (per 100,000 population)

<=0.4

0.41 - 2.2

>2.2

South Dakota, 2011

Primary and secondary syphilis

- CASES: South Dakota reported 0 cases in 2011.
- RATES: South Dakota reported 0 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In South Dakota, the rate among males was 0.0 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.0 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In South Dakota, the race/ethnicity adjusted rates per 100,000 population were 0.0 among whites, 0.0 among blacks, 0.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.


Congenital syphilis

- CASES: South Dakota reported 0 cases in 2011.
- RATES: South Dakota had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, South Dakota, 2002 – 2011



Total P&S Syphilis Rates by County, South Dakota, 2011



Rate (per 100,000 population)



Tennessee, 2011

Primary and secondary syphilis

- CASES: Tennessee reported 278 cases in 2011.
- RATES: Tennessee ranked 18 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 4.4 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Tennessee, the rate among males was 7.9 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 1.0 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Tennessee, the race/ethnicity adjusted rates per 100,000 population were 1.9 among whites, 15.7 among blacks, 2.8 among Hispanics, 0.0 among Asian/Pacific Islanders, and 5.6 among American Indians/Alaska Natives.
- The rate among blacks was 8.3 times that of whites.
- The rate among Hispanics was 1.5 times that of whites.
- The rate among American Indians/Alaska Natives was 2.9 times that of whites.


Congenital syphilis

- CASES: Tennessee reported 8 cases in 2011.
- RATES: Tennessee ranked 11 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 9.4 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Tennessee, 2002 – 2011



Total P&S Syphilis Rates by County, Tennessee, 2011



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Texas, 2011

Primary and secondary syphilis

- CASES: Texas reported 1,169 cases in 2011.
- RATES: Texas ranked 16 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 4.6 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Texas, the rate among males was 7.3 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 2.0 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Texas, the race/ethnicity adjusted rates per 100,000 population were 2.3 among whites, 18.2 among blacks, 3.7 among Hispanics, 1.2 among Asian/Pacific Islanders, and 1.1 among American Indians/Alaska Natives.
- The rate among blacks was 7.9 times that of whites.
- The rate among Hispanics was 1.6 times that of whites.


Congenital syphilis

- CASES: Texas reported 107 cases in 2011.
- RATES: Texas ranked 4 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 26.4 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Texas, 2002 – 2011



Total P&S Syphilis Rates by County, Texas, 2011



Rate (per 100,000 population)



Utah, 2011

Primary and secondary syphilis

- CASES: Utah reported 14 cases in 2011.
- RATES: Utah ranked 49 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 0.5 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Utah, the rate among males was 1.0 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.0 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Utah, the race/ethnicity adjusted rates per 100,000 population were 0.5 among whites, 0.0 among blacks, 0.6 among Hispanics, 0.0 among Asian/Pacific Islanders, and 3.3 among American Indians/Alaska Natives.
- The rate among Hispanics was 1.2 times that of whites.
- The rate among American Indians/Alaska Natives was 6.6 times that of whites.


Congenital syphilis

- CASES: Utah reported 0 cases in 2011.
- RATES: Utah had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Utah, 2002 – 2011



Total P&S Syphilis Rates by County, Utah, 2011



Rate (per 100,000 population)



Vermont, 2011

Primary and secondary syphilis

- CASES: Vermont reported 9 cases in 2011.
- RATES: Vermont ranked 38 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 1.4 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Vermont, the rate among males was 2.9 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.0 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Vermont, the race/ethnicity adjusted rates per 100,000 population were 1.5 among whites, 0.0 among blacks, 0.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.


Congenital syphilis

- CASES: Vermont reported 0 cases in 2011.
- RATES: Vermont had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Vermont, 2002 – 2011



Total P&S Syphilis Rates by County, Vermont, 2011



Rate (per 100,000 population)



Virginia, 2011

Primary and secondary syphilis

- CASES: Virginia reported 213 cases in 2011.
- RATES: Virginia ranked 30 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.7 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Virginia, the rate among males was 5.0 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.4 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Virginia, the race/ethnicity adjusted rates per 100,000 population were 1.2 among whites, 8.2 among blacks, 1.9 among Hispanics, 0.6 among Asian/Pacific Islanders, and 4.2 among American Indians/Alaska Natives.
- The rate among blacks was 6.8 times that of whites.
- The rate among Hispanics was 1.6 times that of whites.
- The rate among American Indians/Alaska Natives was 3.5 times that of whites.


Congenital syphilis

- CASES: Virginia reported 0 cases in 2011.
- RATES: Virginia had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Virginia, 2002 – 2011



Total P&S Syphilis Rates by County, Virginia, 2011



Rate (per 100,000 population)

<=0.4

0.41 - 2.2

>2.2

Washington, 2011

Primary and secondary syphilis

- CASES: Washington reported 328 cases in 2011.
- RATES: Washington ranked 13 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 4.9 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Washington, the rate among males was 9.6 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.2 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Washington, the race/ethnicity adjusted rates per 100,000 population were 3.7 among whites, 9.5 among blacks, 4.2 among Hispanics, 2.1 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 2.6 times that of whites.
- The rate among Hispanics was 1.1 times that of whites.


Congenital syphilis

- CASES: Washington reported 2 cases in 2011.
- RATES: Washington ranked 24 among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 2.2 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Washington, 2002 – 2011



Total P&S Syphilis Rates by County, Washington, 2011



Rate (per 100,000 population)



West Virginia, 2011

Primary and secondary syphilis

- CASES: West Virginia reported 4 cases in 2011.
- RATES: West Virginia ranked 50 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 0.2 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In West Virginia, the rate among males was 0.4 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.0 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In West Virginia, the race/ethnicity adjusted rates per 100,000 population were 0.2 among whites, 0.0 among blacks, 0.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.


Congenital syphilis

- CASES: West Virginia reported 0 cases in 2011.
- RATES: West Virginia had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, West Virginia, 2002 – 2011



Total P&S Syphilis Rates by County, West Virginia, 2011



Rate (per 100,000 population)

≤0.4 0.41 - 2.2 >2.2

Wisconsin, 2011

Primary and secondary syphilis

- CASES: Wisconsin reported 65 cases in 2011.
- RATES: Wisconsin ranked 40 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 1.1 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Wisconsin, the rate among males was 2.1 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.2 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Wisconsin, the race/ethnicity adjusted rates per 100,000 population were 0.6 among whites, 7.7 among blacks, 1.2 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 12.8 times that of whites.
- The rate among Hispanics was 2 times that of whites.


Congenital syphilis

- CASES: Wisconsin reported 0 cases in 2011.
- RATES: Wisconsin had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Wisconsin, 2002 – 2011



Total P&S Syphilis Rates by County, Wisconsin, 2011



Rate (per 100,000 population)

≤0.4 0.41 - 2.2 >2.2

Wyoming, 2011

Primary and secondary syphilis

- CASES: Wyoming reported 0 cases in 2011.
- RATES: Wyoming reported 0 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Wyoming, the rate among males was 0.0 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.0 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Wyoming, the race/ethnicity adjusted rates per 100,000 population were 0.0 among whites, 0.0 among blacks, 0.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.


Congenital syphilis

- CASES: Wyoming reported 0 cases in 2011.
- RATES: Wyoming had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Wyoming, 2002 – 2011



Total P&S Syphilis Rates by County, Wyoming, 2011



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Guam, 2011

Primary and secondary syphilis

- CASES: Guam reported 5 cases in 2011.
- RATES: Guam ranked 28 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 2.8 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex


- In Guam, the rate among males was 3.3 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 2.2 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.

Rates of primary and secondary syphilis, by race/ethnicity


Congenital syphilis

- CASES: Guam reported 0 cases in 2011.
- RATES: Guam had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Guam, 2002 – 2011



Total P&S Syphilis Rate, Guam, 2011



Puerto Rico, 2011

Primary and secondary syphilis

- CASES: Puerto Rico reported 254 cases in 2011.
- RATES: Puerto Rico ranked 6 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 6.8 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex


- In Puerto Rico, the rate among males was 13.3 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.9 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.

Rates of primary and secondary syphilis, by race/ethnicity


Congenital syphilis

- CASES: Puerto Rico reported 2 cases in 2011.
- RATES: Puerto Rico ranked 19 among among 25 areas (23 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 4.3 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Puerto Rico, 2002 – 2011



Total P&S Syphilis Rates by Municipality, Puerto Rico, 2011



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2

Virgin Islands, 2011

Primary and secondary syphilis

- CASES: Virgin Islands reported 0 cases in 2011.
- RATES: Virgin Islands reported 0 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex


- In Virgin Islands, the rate among males was 0.0 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.0 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.

Rates of primary and secondary syphilis, by race/ethnicity

Congenital syphilis

- CASES: Virgin Islands reported 0 cases in 2011.
- RATES: Virgin Islands had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Virgin Islands, 2002 – 2011



Total P&S Syphilis Rate, Virgin Islands, 2011



Rate (per 100,000 population)

≤0.4

0.41 - 2.2

>2.2