

Wisconsin, 2011

Primary and secondary syphilis

- CASES: Wisconsin reported 65 cases in 2011.
- RATES: Wisconsin ranked 40 among 51 areas (48 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 1.1 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Wisconsin, the rate among males was 2.1 per 100,000 population compared to the U.S. male rate of 8.2 per 100,000.
- The rate among females was 0.2 per 100,000 compared to the U.S. female rate of 1.0 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Wisconsin, the race/ethnicity adjusted rates per 100,000 population were 0.6 among whites, 7.7 among blacks, 1.2 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 12.8 times that of whites.
- The rate among Hispanics was 2 times that of whites.

Congenital syphilis

- CASES: Wisconsin reported 0 cases in 2011.
- RATES: Wisconsin had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Wisconsin, 2002 – 2011



Total P&S Syphilis Rates by County, Wisconsin, 2011

