

District of Columbia, 2012

Primary and secondary syphilis

- **CASES:** District of Columbia reported 165 cases in 2012.
- **RATES:** District of Columbia reported 26.7 cases per 100,000 population compared to the U.S. rate of 5.0.

Rates of primary and secondary syphilis, by sex

- In District of Columbia, the rate among males was 54.4 cases per 100,000 population compared to the U.S. male rate of 9.3.
- The rate among females was 1.8 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In District of Columbia, the race/ethnicity adjusted rates per 100,000 population were 25.7 among whites, 29.3 among blacks, 22.1 among Hispanics, 0.0 among Asians, 0.0 among Native Hawaiians and Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 1.1 times that of whites.


Congenital syphilis

- **CASES:** District of Columbia reported 0 cases in 2012.
- **RATES:** District of Columbia had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 7.8.

Syphilis: P&S Rates and Congenital Cases, District of Columbia, 2003–2012



Total P&S Syphilis Rate, District of Columbia, 2012



Rate (per 100,000 population)

