

Alaska, 2013

Primary and secondary syphilis

- **CASES:** Alaska reported 23 cases in 2013.
- **RATES:** Alaska ranked 32 among 50 states reporting at least one case of P&S syphilis with 3.1 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Alaska, the rate among males was 5.5 per 100,000 population compared to the U.S. male rate of 10.3 cases per 100,000.
- The rate among females was 0.6 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity*

- In Alaska, the race/ethnicity adjusted rates per 100,000 population were 1.7 among whites, 9.4 among blacks, 2.2 among Hispanics, 3.8 among Asian/Pacific Islanders, and 7.7 among American Indians/Alaska Natives.
- The rate among blacks was 5.5 times that of whites.
- The rate among Hispanics was 1.3 times that of whites.
- The rate among Asian/Pacific Islanders was 2.2 times that of whites.
- The rate among American Indians/Alaska Natives was 4.5 times that of whites.


Congenital syphilis

- **CASES:** Alaska reported 1 case in 2013.
- **RATES:** Alaska ranked 12 among 25 states reporting at least one case of congenital syphilis with a rate of 8.7 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Alaska, 2004–2013



Total P&S Syphilis Rates by County, Alaska, 2013



Rate (per 100,000 population)



*OMB-compliant race values not available. NCHS bridged population used to calculate rates.