

Alabama, 2013

Primary and secondary syphilis

- CASES: Alabama reported 183 cases in 2013.
- RATES: Alabama ranked 23 among 50 states reporting at least one case of P&S syphilis with 3.8 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Alabama, the rate among males was 6.9 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.9 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Alabama, the race/ethnicity adjusted rates per 100,000 population were 1.3 among whites, 9.1 among blacks, 1.0 among Hispanics, 0.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 6.8 times that of whites.


Congenital syphilis

- CASES: Alabama reported 2 cases in 2013.
- RATES: Alabama ranked 20 among 25 states reporting at least one case of congenital syphilis with a rate of 3.3 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Alabama, 2004–2013



Total P&S Syphilis Rates by County, Alabama, 2013



Rate (per 100,000 population)

