

District of Columbia, 2013

Primary and secondary syphilis

- **CASES:** District of Columbia reported 168 cases in 2013.
- **RATES:** District of Columbia reported 26.6 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In District of Columbia, the rate among males was 49.8 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 5.7 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In District of Columbia, the race/ethnicity adjusted rates per 100,000 population were 18.7 among whites, 33.9 among blacks, 9.6 among Hispanics, 12.9 among Asians, 266 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 1.8 times that of whites.
- The rate among Native Hawaiian/Other Pacific Islanders was 14.2 times that of whites.


Congenital syphilis

- **CASES:** District of Columbia reported 2 cases in 2013.
- **RATES:** District of Columbia had a rate of 21.8 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, District of Columbia, 2004–2013



Total P&S Syphilis Rate, District of Columbia, 2013



Rate (per 100,000 population)

