

Hawaii, 2013

Primary and secondary syphilis

- CASES: Hawaii reported 46 cases in 2013.
- RATES: Hawaii ranked 30 among 50 states reporting at least one case of P&S syphilis with 3.3 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Hawaii, the rate among males was 6.6 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.0 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Hawaii, the race/ethnicity adjusted rates per 100,000 population were 3.8 among whites, 7.5 among blacks, 3.0 among Hispanics, 2.5 among Asians, 3.8 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 2 times that of whites.

Congenital syphilis

- CASES: Hawaii reported 0 cases in 2013.
- RATES: Hawaii had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Hawaii, 2004–2013



Total P&S Syphilis Rates by County, Hawaii, 2013

