

Iowa, 2013

Primary and secondary syphilis

- CASES: Iowa reported 106 cases in 2013.
- RATES: Iowa ranked 28 among 50 states reporting at least one case of P&S syphilis with 3.4 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Iowa, the rate among males was 6.3 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.6 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Iowa, the race/ethnicity adjusted rates per 100,000 population were 2.7 among whites, 19.4 among blacks, 2.5 among Hispanics, 5.1 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 11.4 among American Indians/Alaska Natives.
- The rate among blacks was 7.1 times that of whites.
- The rate among Asians was 1.9 times that of whites.
- The rate among American Indians/Alaska Natives was 4.2 times that of whites.


Congenital syphilis

- CASES: Iowa reported 0 cases in 2013.
- RATES: Iowa had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Iowa, 2004–2013



Total P&S Syphilis Rates by County, Iowa, 2013



Rate (per 100,000 population)

