

Indiana, 2013

Primary and secondary syphilis

- CASES: Indiana reported 215 cases in 2013.
- RATES: Indiana ranked 31 among 50 states reporting at least one case of P&S syphilis with 3.3 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Indiana, the rate among males was 6.1 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.5 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Indiana, the race/ethnicity adjusted rates per 100,000 population were 1.8 among whites, 14.9 among blacks, 5.6 among Hispanics, 1.8 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 8.5 times that of whites.
- The rate among Hispanics was 3.2 times that of whites.


Congenital syphilis

- CASES: Indiana reported 0 cases in 2013.
- RATES: Indiana had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Indiana, 2004–2013



Total P&S Syphilis Rates by County, Indiana, 2013



Rate (per 100,000 population)

