

# Kentucky, 2013

## Primary and secondary syphilis

- **CASES:** Kentucky reported 122 cases in 2013.
- **RATES:** Kentucky ranked 35 among 50 states reporting at least one case of P&S syphilis with 2.8 cases per 100,000 population compared to the U.S. rate of 5.5.

## Rates of primary and secondary syphilis, by sex

- In Kentucky, the rate among males was 4.9 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.8 compared to the U.S. female rate of 0.9.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Kentucky, the race/ethnicity adjusted rates per 100,000 population were 1.6 among whites, 14.8 among blacks, 4.9 among Hispanics, 1.8 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 9 times that of whites.
- The rate among Hispanics was 3 times that of whites.
- The rate among Asians was 1.1 times that of whites.


## Congenital syphilis

- **CASES:** Kentucky reported 4 cases in 2013.
- **RATES:** Kentucky ranked 15 among 25 states reporting at least one case of congenital syphilis with a rate of 7.2 cases per 100,000 live births compared to the U.S. rate of 8.7.

## Syphilis: P&S Rates and Congenital Cases, Kentucky, 2004–2013



## Total P&S Syphilis Rates by County, Kentucky, 2013



Rate (per 100,000 population)

