

Massachusetts, 2013

Primary and secondary syphilis

- CASES: Massachusetts reported 360 cases in 2013.
- RATES: Massachusetts ranked 14 among 50 states reporting at least one case of P&S syphilis with 5.4 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Massachusetts, the rate among males was 10.7 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.5 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Massachusetts, the race/ethnicity adjusted rates per 100,000 population were 4.0 among whites, 11.6 among blacks, 10.1 among Hispanics, 2.4 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 33.6 among American Indians/Alaska Natives.
- The rate among blacks was 2.9 times that of whites.
- The rate among Hispanics was 2.5 times that of whites.
- The rate among American Indians/Alaska Natives was 8.4 times that of whites.


Congenital syphilis

- CASES: Massachusetts reported 4 cases in 2013.
- RATES: Massachusetts ranked 17 among 25 states reporting at least one case of congenital syphilis with a rate of 5.5 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Massachusetts, 2004–2013



Total P&S Syphilis Rates by County, Massachusetts, 2013



Rate (per 100,000 population)

