

Michigan, 2013

Primary and secondary syphilis

- CASES: Michigan reported 487 cases in 2013.
- RATES: Michigan ranked 16 among 50 states reporting at least one case of P&S syphilis with 4.9 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Michigan, the rate among males was 9.4 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.6 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Michigan, the race/ethnicity adjusted rates per 100,000 population were 1.7 among whites, 24.0 among blacks, 3.5 among Hispanics, 0.8 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 1.8 among American Indians/Alaska Natives.
- The rate among blacks was 13.9 times that of whites.
- The rate among Hispanics was 2 times that of whites.


Congenital syphilis

- CASES: Michigan reported 9 cases in 2013.
- RATES: Michigan ranked 13 among among 25 states reporting at least one case of congenital syphilis with a rate of 7.9 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Michigan, 2004–2013



Total P&S Syphilis Rates by County, Michigan, 2013



Rate (per 100,000 population)

