

# Minnesota, 2013

## Primary and secondary syphilis

- CASES: Minnesota reported 193 cases in 2013.
- RATES: Minnesota ranked 27 among 50 states reporting at least one case of P&S syphilis with 3.6 cases per 100,000 population compared to the U.S. rate of 5.5.

## Rates of primary and secondary syphilis, by sex

- In Minnesota, the rate among males was 6.7 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.4 compared to the U.S. female rate of 0.9.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Minnesota, the race/ethnicity adjusted rates per 100,000 population were 2.5 among whites, 16.4 among blacks, 3.4 among Hispanics, 3.0 among Asians, 46.7 among Native Hawaiian/Other Pacific Islanders, and 3.5 among American Indians/Alaska Natives.
- The rate among blacks was 6.6 times that of whites.
- The rate among Hispanics was 1.4 times that of whites.
- The rate among Asians was 1.2 times that of whites.
- The rate among Native Hawaiian/Other Pacific Islanders was 19 times that of whites.
- The rate among American Indians/Alaska Natives was 1.4 times that of whites.


## Congenital syphilis

- CASES: Minnesota reported 0 cases in 2013.
- RATES: Minnesota had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

## Syphilis: P&S Rates and Congenital Cases, Minnesota, 2004–2013



## Total P&S Syphilis Rates by County, Minnesota, 2013



Rate (per 100,000 population)

≤0.4
  0.41 - 2.2
  >2.2