

Missouri, 2013

Primary and secondary syphilis

- CASES: Missouri reported 251 cases in 2013.
- RATES: Missouri ranked 19 among 50 states reporting at least one case of P&S syphilis with 4.2 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Missouri, the rate among males was 7.9 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.6 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Missouri, the race/ethnicity adjusted rates per 100,000 population were 2.0 among whites, 18.4 among blacks, 6.7 among Hispanics, 0.0 among Asians, 31.2 among Native Hawaiian/Other Pacific Islanders, and 8.1 among American Indians/Alaska Natives.
- The rate among blacks was 9.2 times that of whites.
- The rate among Hispanics was 3.3 times that of whites.
- The rate among Native Hawaiian/Other Pacific Islanders was 15.6 times that of whites.
- The rate among American Indians/Alaska Natives was 4 times that of whites.


Congenital syphilis

- CASES: Missouri reported 3 cases in 2013.
- RATES: Missouri ranked 19 among among 25 states reporting at least one case of congenital syphilis with a rate of 3.9 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Missouri, 2004–2013



Total P&S Syphilis Rates by County, Missouri, 2013



Rate (per 100,000 population)

