

North Carolina, 2013

Primary and secondary syphilis

- CASES: North Carolina reported 404 cases in 2013.
- RATES: North Carolina ranked 20 among 50 states reporting at least one case of P&S syphilis with 4.1 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In North Carolina, the rate among males was 7.7 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.7 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In North Carolina, the race/ethnicity adjusted rates per 100,000 population were 1.6 among whites, 13.5 among blacks, 1.3 among Hispanics, 0.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 3.6 among American Indians/Alaska Natives.
- The rate among blacks was 8.5 times that of whites.
- The rate among American Indians/Alaska Natives was 2.3 times that of whites.


Congenital syphilis

- CASES: North Carolina reported 1 case in 2013.
- RATES: North Carolina ranked 25 among 25 states reporting at least one case of congenital syphilis with a rate of 0.8 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, North Carolina, 2004–2013



Total P&S Syphilis Rates by County, North Carolina, 2013



Rate (per 100,000 population)

