

New Jersey, 2013

Primary and secondary syphilis

- CASES: New Jersey reported 233 cases in 2013.
- RATES: New Jersey ranked 36 among 50 states reporting at least one case of P&S syphilis with 2.6 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In New Jersey, the rate among males was 5.1 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.3 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In New Jersey, the race/ethnicity adjusted rates per 100,000 population were 1.4 among whites, 7.8 among blacks, 4.0 among Hispanics, 0.4 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 5.5 times that of whites.
- The rate among Hispanics was 2.8 times that of whites.


Congenital syphilis

- CASES: New Jersey reported 0 cases in 2013.
- RATES: New Jersey had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, New Jersey, 2004–2013



Total P&S Syphilis Rates by County, New Jersey, 2013



Rate (per 100,000 population)

