

New Mexico, 2013

Primary and secondary syphilis

- CASES: New Mexico reported 78 cases in 2013.
- RATES: New Mexico ranked 25 among 50 states reporting at least one case of P&S syphilis with 3.7 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In New Mexico, the rate among males was 5.6 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 1.9 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In New Mexico, the race/ethnicity adjusted rates per 100,000 population were 2.0 among whites, 16.1 among blacks, 3.8 among Hispanics, 0.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 3.4 among American Indians/Alaska Natives.
- The rate among blacks was 7.9 times that of whites.
- The rate among Hispanics was 1.8 times that of whites.
- The rate among American Indians/Alaska Natives was 1.6 times that of whites.


Congenital syphilis

- CASES: New Mexico reported 2 cases in 2013.
- RATES: New Mexico ranked 14 among 25 states reporting at least one case of congenital syphilis with a rate of 7.2 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, New Mexico, 2004–2013



Total P&S Syphilis Rates by County, New Mexico, 2013



Rate (per 100,000 population)

