

New York, 2013

Primary and secondary syphilis

- CASES: New York reported 1,459 cases in 2013.
- RATES: New York ranked 6 among 50 states reporting at least one case of P&S syphilis with 7.5 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In New York, the rate among males was 14.8 per 100,000 population compared to the U.S. male rate of 10.3 cases per 100,000.
- The rate among females was 0.4 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity*

- In New York, the race/ethnicity adjusted rates per 100,000 population were 3.7 among whites, 15.8 among blacks, 10.0 among Hispanics, 2.9 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 4.3 times that of whites.
- The rate among Hispanics was 2.7 times that of whites.


Congenital syphilis

- CASES: New York reported 11 cases in 2013.
- RATES: New York ranked 18 among 25 states reporting at least one case of congenital syphilis with a rate of 4.5 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, New York, 2004–2013



Total P&S Syphilis Rates by County, New York, 2013



Rate (per 100,000 population)



*OMB-compliant race values not available. NCHS bridged population used to calculate rates.