

Ohio, 2013

Primary and secondary syphilis

- CASES: Ohio reported 436 cases in 2013.
- RATES: Ohio ranked 24 among 50 states reporting at least one case of P&S syphilis with 3.8 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Ohio, the rate among males was 6.6 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 1.1 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Ohio, the race/ethnicity adjusted rates per 100,000 population were 2.1 among whites, 15.1 among blacks, 4.2 among Hispanics, 1.0 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 7.3 times that of whites.
- The rate among Hispanics was 2 times that of whites.


Congenital syphilis

- CASES: Ohio reported 17 cases in 2013.
- RATES: Ohio ranked 9 among 25 states reporting at least one case of congenital syphilis with a rate of 12.2 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Ohio, 2004–2013



Total P&S Syphilis Rates by County, Ohio, 2013



Rate (per 100,000 population)

