

# Pennsylvania, 2013

## Primary and secondary syphilis

- CASES: Pennsylvania reported 471 cases in 2013.
- RATES: Pennsylvania ranked 26 among 50 states reporting at least one case of P&S syphilis with 3.7 cases per 100,000 population compared to the U.S. rate of 5.5.

## Rates of primary and secondary syphilis, by sex

- In Pennsylvania, the rate among males was 7.1 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.4 compared to the U.S. female rate of 0.9.


## Rates of primary and secondary syphilis, by race/ethnicity

- In Pennsylvania, the race/ethnicity adjusted rates per 100,000 population were 1.7 among whites, 16.7 among blacks, 5.0 among Hispanics, 0.8 among Asians, 31.4 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 10 times that of whites.
- The rate among Hispanics was 3 times that of whites.
- The rate among Native Hawaiian/Other Pacific Islanders was 18.7 times that of whites.


## Congenital syphilis

- CASES: Pennsylvania reported 2 cases in 2013.
- RATES: Pennsylvania ranked 24 among 25 states reporting at least one case of congenital syphilis with a rate of 1.4 cases per 100,000 live births compared to the U.S. rate of 8.7.

## Syphilis: P&S Rates and Congenital Cases, Pennsylvania, 2004–2013



## Total P&S Syphilis Rates by County, Pennsylvania, 2013



Rate (per 100,000 population)

