

Rhode Island, 2013

Primary and secondary syphilis

- CASES: Rhode Island reported 45 cases in 2013.
- RATES: Rhode Island ranked 18 among 50 states reporting at least one case of P&S syphilis with 4.3 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Rhode Island, the rate among males was 8.7 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.2 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Rhode Island, the race/ethnicity adjusted rates per 100,000 population were 3.8 among whites, 7.0 among blacks, 4.3 among Hispanics, 9.2 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 1.8 times that of whites.
- The rate among Hispanics was 1.2 times that of whites.
- The rate among Asians was 2.4 times that of whites.

Congenital syphilis

- CASES: Rhode Island reported 0 cases in 2013.
- RATES: Rhode Island had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Rhode Island, 2004–2013



Total P&S Syphilis Rates by County, Rhode Island, 2013



Rate (per 100,000 population)

