

South Carolina, 2013

Primary and secondary syphilis

- CASES: South Carolina reported 271 cases in 2013.
- RATES: South Carolina ranked 11 among 50 states reporting at least one case of P&S syphilis with 5.7 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In South Carolina, the rate among males was 10.1 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 1.6 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In South Carolina, the race/ethnicity adjusted rates per 100,000 population were 2.0 among whites, 15.8 among blacks, 1.6 among Hispanics, 1.5 among Asians, 0.0 among Native Hawaiian/Other Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 7.8 times that of whites.


Congenital syphilis

- CASES: South Carolina reported 1 case in 2013.
- RATES: South Carolina ranked 23 among 25 states reporting at least one case of congenital syphilis with a rate of 1.7 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, South Carolina, 2004–2013



Total P&S Syphilis Rates by County, South Carolina, 2013



Rate (per 100,000 population)

≤0.4
 0.41 - 2.2
 >2.2