

Washington, 2013

Primary and secondary syphilis

- CASES: Washington reported 284 cases in 2013.
- RATES: Washington ranked 21 among 50 states reporting at least one case of P&S syphilis with 4.1 cases per 100,000 population compared to the U.S. rate of 5.5.

Rates of primary and secondary syphilis, by sex

- In Washington, the rate among males was 7.9 cases per 100,000 population compared to the U.S. male rate of 10.3.
- The rate among females was 0.4 compared to the U.S. female rate of 0.9.


Rates of primary and secondary syphilis, by race/ethnicity

- In Washington, the race/ethnicity adjusted rates per 100,000 population were 3.9 among whites, 12.2 among blacks, 4.8 among Hispanics, 1.2 among Asians, 2.4 among Native Hawaiian/Other Pacific Islanders, and 4.4 among American Indians/Alaska Natives.
- The rate among blacks was 3.1 times that of whites.
- The rate among Hispanics was 1.2 times that of whites.
- The rate among American Indians/Alaska Natives was 1.1 times that of whites.


Congenital syphilis

- CASES: Washington reported 0 cases in 2013.
- RATES: Washington had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7.

Syphilis: P&S Rates and Congenital Cases, Washington, 2004–2013



Total P&S Syphilis Rates by County, Washington, 2013



Rate (per 100,000 population)

